

KLE axioms

KLE001+0.ax Idempotent semirings

$\forall a, b: a + b = b + a \quad \text{fof(additive_commutativity, axiom)}$
 $\forall c, b, a: a + (b + c) = (a + b) + c \quad \text{fof(additive_associativity, axiom)}$
 $\forall a: a + 0 = a \quad \text{fof(additive_identity, axiom)}$
 $\forall a: a + a = a \quad \text{fof(additive_idempotence, axiom)}$
 $\forall a, b, c: a \cdot (b \cdot c) = (a \cdot b) \cdot c \quad \text{fof(multiplicative_associativity, axiom)}$
 $\forall a: a \cdot 1 = a \quad \text{fof(multiplicative_right_identity, axiom)}$
 $\forall a: 1 \cdot a = a \quad \text{fof(multiplicative_left_identity, axiom)}$
 $\forall a, b, c: a \cdot (b + c) = a \cdot b + a \cdot c \quad \text{fof(right_distributivity, axiom)}$
 $\forall a, b, c: (a + b) \cdot c = a \cdot c + b \cdot c \quad \text{fof(left_distributivity, axiom)}$
 $\forall a: a \cdot 0 = 0 \quad \text{fof(right_annihilation, axiom)}$
 $\forall a: 0 \cdot a = 0 \quad \text{fof(left_annihilation, axiom)}$
 $\forall a, b: (a \leq b \iff a + b = b) \quad \text{fof(order, axiom)}$

KLE001+1.ax Characterisation of tests by complement predicate

$\forall x_0: (\text{test}(x_0) \iff \exists x_1: x'_1 = x_0) \quad \text{fof(test}_1, \text{axiom})$
 $\forall x_0, x_1: (x'_1 = x_0 \iff (x_0 \cdot x_1 = 0 \text{ and } x_1 \cdot x_0 = 0 \text{ and } x_0 + x_1 = 1)) \quad \text{fof(test}_2, \text{axiom})$
 $\forall x_0, x_1: (\text{test}(x_0) \Rightarrow (c(x_0) = x_1 \iff x'_0 = x_1)) \quad \text{fof(test}_3, \text{axiom})$
 $\forall x_0: (\neg \text{test}(x_0) \Rightarrow c(x_0) = 0) \quad \text{fof(test}_4, \text{axiom})$

KLE001+2.ax de Morgan's laws for tests

$\forall x_0, x_1: ((\text{test}(x_0) \text{ and } \text{test}(x_1)) \Rightarrow c(x_0 + x_1) = c(x_0) \cdot c(x_1)) \quad \text{fof(test_deMorgan}_1, \text{axiom})$
 $\forall x_0, x_1: ((\text{test}(x_0) \text{ and } \text{test}(x_1)) \Rightarrow c(x_0 \cdot x_1) = c(x_0) + c(x_1)) \quad \text{fof(test_deMorgan}_2, \text{axiom})$

KLE001+3.ax Universal characterisation of meet

$\forall x_0, x_1, x_2: (\text{ismeet}(x_2, x_0, x_1) \iff (x_2 \leq x_0 \text{ and } x_2 \leq x_1 \text{ and } \forall x_3: ((x_3 \leq x_0 \text{ and } x_3 \leq x_1) \Rightarrow x_3 \leq x_2))) \quad \text{fof(ismeet, axiom)}$
 $\forall x_0, x_1, x_2: (\text{ismeetu}(x_2, x_0, x_1) \iff \forall x_3: ((x_3 \leq x_0 \text{ and } x_3 \leq x_1) \iff x_3 \leq x_2)) \quad \text{fof(ismeetu, axiom)}$

KLE001+4.ax Boolean domain, antidomain, codomain, coantidomain

$\forall x_0: \text{ad}(x_0) \cdot x_0 = 0 \quad \text{fof(domain}_1, \text{axiom})$
 $\forall x_0, x_1: \text{ad}(x_0 \cdot x_1) + \text{ad}(x_0 \cdot \text{ad}(\text{ad}(x_1))) = \text{ad}(x_0 \cdot \text{ad}(\text{ad}(x_1))) \quad \text{fof(domain}_2, \text{axiom})$
 $\forall x_0: \text{ad}(\text{ad}(x_0)) + \text{ad}(x_0) = 1 \quad \text{fof(domain}_3, \text{axiom})$
 $\forall x_0: \text{dom}(x_0) = \text{ad}(\text{ad}(x_0)) \quad \text{fof(domain}_4, \text{axiom})$
 $\forall x_0: x_0 \cdot \text{coad}(x_0) = 0 \quad \text{fof(codomain}_1, \text{axiom})$
 $\forall x_0, x_1: \text{coad}(x_0 \cdot x_1) + \text{coad}(\text{coad}(\text{coad}(x_0)) \cdot x_1) = \text{coad}(\text{coad}(\text{coad}(x_0)) \cdot x_1) \quad \text{fof(codomain}_2, \text{axiom})$
 $\forall x_0: \text{coad}(\text{coad}(x_0)) + \text{coad}(x_0) = 1 \quad \text{fof(codomain}_3, \text{axiom})$
 $\forall x_0: \text{cod}(x_0) = \text{coad}(\text{coad}(x_0)) \quad \text{fof(codomain}_4, \text{axiom})$

KLE001+5.ax Domain (not Boolean domain!)

$\forall x_0: x_0 + \text{dom}(x_0) \cdot x_0 = \text{dom}(x_0) \cdot x_0 \quad \text{fof(domain}_1, \text{axiom})$
 $\forall x_0, x_1: \text{dom}(x_0 \cdot x_1) = \text{dom}(x_0 \cdot \text{dom}(x_1)) \quad \text{fof(domain}_2, \text{axiom})$
 $\forall x_0: \text{dom}(x_0) + 1 = 1 \quad \text{fof(domain}_3, \text{axiom})$
 $\text{dom}(0) = 0 \quad \text{fof(domain}_4, \text{axiom})$
 $\forall x_0, x_1: \text{dom}(x_0 + x_1) = \text{dom}(x_0) + \text{dom}(x_1) \quad \text{fof(domain}_5, \text{axiom})$

KLE001+6.ax Modal operators

$\forall x_0: c(x_0) = \text{ad}(\text{dom}(x_0)) \quad \text{fof(complement, axiom)}$
 $\forall x_0, x_1: \text{domain_difference}(x_0, x_1) = \text{dom}(x_0) \cdot \text{ad}(x_1) \quad \text{fof(domain_difference, axiom)}$
 $\forall x_0, x_1: \text{forward_diamond}(x_0, x_1) = \text{dom}(x_0 \cdot \text{dom}(x_1)) \quad \text{fof(forward_diamond, axiom)}$
 $\forall x_0, x_1: \text{backward_diamond}(x_0, x_1) = \text{cod}(\text{cod}(x_1) \cdot x_0) \quad \text{fof(backward_diamond, axiom)}$
 $\forall x_0, x_1: \text{forward_box}(x_0, x_1) = c(\text{forward_diamond}(x_0, c(x_1))) \quad \text{fof(forward_box, axiom)}$
 $\forall x_0, x_1: \text{backward_box}(x_0, x_1) = c(\text{backward_diamond}(x_0, c(x_1))) \quad \text{fof(backward_box, axiom)}$

KLE001+7.ax Divergence Kleene algebras

$\forall x_0: \text{forward_diamond}(x_0, \text{divergence}(x_0)) = \text{divergence}(x_0) \quad \text{fof(divergence}_1, \text{axiom})$
 $\forall x_0, x_1, x_2: (\text{dom}(x_0) + (\text{forward_diamond}(x_1, \text{dom}(x_0)) + \text{dom}(x_2)) = \text{forward_diamond}(x_1, \text{dom}(x_0)) + \text{dom}(x_2) \Rightarrow \text{dom}(x_0) + (\text{divergence}(x_1) + \text{forward_diamond}(x_1^*, \text{dom}(x_2))) = \text{divergence}(x_1) + \text{forward_diamond}(x_1^*, \text{dom}(x_2))) \quad \text{fof(divergence}_2, \text{axiom})$

KLE002+0.ax Kleene algebra

$\forall a, b: a + b = b + a \quad \text{fof(additive_commutativity, axiom)}$
 $\forall c, b, a: a + (b + c) = (a + b) + c \quad \text{fof(additive_associativity, axiom)}$

$\forall a: a + 0 = a \quad \text{fof(additive_identity, axiom)}$
 $\forall a: a + a = a \quad \text{fof(additive_idempotence, axiom)}$
 $\forall a, b, c: a \cdot (b \cdot c) = (a \cdot b) \cdot c \quad \text{fof(multiplicative_associativity, axiom)}$
 $\forall a: a \cdot 1 = a \quad \text{fof(multiplicative_right_identity, axiom)}$
 $\forall a: 1 \cdot a = a \quad \text{fof(multiplicative_left_identity, axiom)}$
 $\forall a, b, c: a \cdot (b + c) = a \cdot b + a \cdot c \quad \text{fof(right_distributivity, axiom)}$
 $\forall a, b, c: (a + b) \cdot c = a \cdot c + b \cdot c \quad \text{fof(left_distributivity, axiom)}$
 $\forall a: a \cdot 0 = 0 \quad \text{fof(right_annihilation, axiom)}$
 $\forall a: 0 \cdot a = 0 \quad \text{fof(left_annihilation, axiom)}$
 $\forall a, b: (a \leq b \iff a + b = b) \quad \text{fof(order, axiom)}$
 $\forall a: 1 + a \cdot a^* \leq a^* \quad \text{fof(star_unfold_right, axiom)}$
 $\forall a: 1 + a^* \cdot a \leq a^* \quad \text{fof(star_unfold_left, axiom)}$
 $\forall a, b, c: (a \cdot b + c \leq b \Rightarrow a^* \cdot c \leq b) \quad \text{fof(star_induction_left, axiom)}$
 $\forall a, b, c: (a \cdot b + c \leq a \Rightarrow c \cdot b^* \leq a) \quad \text{fof(star_induction_right, axiom)}$

KLE003+0.ax Omega algebra

$\forall a, b: a + b = b + a \quad \text{fof(additive_commutativity, axiom)}$
 $\forall c, b, a: a + (b + c) = (a + b) + c \quad \text{fof(additive_associativity, axiom)}$
 $\forall a: a + 0 = a \quad \text{fof(additive_identity, axiom)}$
 $\forall a: a + a = a \quad \text{fof(additive_idempotence, axiom)}$
 $\forall a, b, c: a \cdot (b \cdot c) = (a \cdot b) \cdot c \quad \text{fof(multiplicative_associativity, axiom)}$
 $\forall a: a \cdot 1 = a \quad \text{fof(multiplicative_right_identity, axiom)}$
 $\forall a: 1 \cdot a = a \quad \text{fof(multiplicative_left_identity, axiom)}$
 $\forall a, b, c: a \cdot (b + c) = a \cdot b + a \cdot c \quad \text{fof(right_distributivity, axiom)}$
 $\forall a, b, c: (a + b) \cdot c = a \cdot c + b \cdot c \quad \text{fof(left_distributivity, axiom)}$
 $\forall a: a \cdot 0 = 0 \quad \text{fof(right_annihilation, axiom)}$
 $\forall a: 0 \cdot a = 0 \quad \text{fof(left_annihilation, axiom)}$
 $\forall a, b: (a \leq b \iff a + b = b) \quad \text{fof(order, axiom)}$
 $\forall a: 1 + a \cdot a^* \leq a^* \quad \text{fof(star_unfold_right, axiom)}$
 $\forall a: 1 + a^* \cdot a \leq a^* \quad \text{fof(star_unfold_left, axiom)}$
 $\forall a, b, c: (a \cdot b + c \leq b \Rightarrow a^* \cdot c \leq b) \quad \text{fof(star_induction_left, axiom)}$
 $\forall a, b, c: (a \cdot b + c \leq a \Rightarrow c \cdot b^* \leq a) \quad \text{fof(star_induction_right, axiom)}$
 $\forall a: a \cdot a^\omega = a^\omega \quad \text{fof(omega_unfold, axiom)}$
 $\forall a, b, c: (a \leq b \cdot a + c \Rightarrow a \leq b^\omega + b^* \cdot c) \quad \text{fof(omega_co_induction, axiom)}$

KLE004+0.ax Demonic Refinement Algebra

$\forall a, b: a + b = b + a \quad \text{fof(additive_commutativity, axiom)}$
 $\forall c, b, a: a + (b + c) = (a + b) + c \quad \text{fof(additive_associativity, axiom)}$
 $\forall a: a + 0 = a \quad \text{fof(additive_identity, axiom)}$
 $\forall a: a + a = a \quad \text{fof(idempotence, axiom)}$
 $\forall a, b, c: a \cdot (b \cdot c) = (a \cdot b) \cdot c \quad \text{fof(multiplicative_associativity, axiom)}$
 $\forall a: a \cdot 1 = a \quad \text{fof(multiplicative_right_identity, axiom)}$
 $\forall a: 1 \cdot a = a \quad \text{fof(multiplicative_left_identity, axiom)}$
 $\forall a, b, c: a \cdot (b + c) = a \cdot b + a \cdot c \quad \text{fof(distributivity}_1\text{, axiom)}$
 $\forall a, b, c: (a + b) \cdot c = a \cdot c + b \cdot c \quad \text{fof(distributivity}_2\text{, axiom)}$
 $\forall a: 0 \cdot a = 0 \quad \text{fof(left_annihilation, axiom)}$
 $\forall a: 1 + a \cdot a^* = a^* \quad \text{fof(star_unfold}_1\text{, axiom)}$
 $\forall a: 1 + a^* \cdot a = a^* \quad \text{fof(star_unfold}_2\text{, axiom)}$
 $\forall a, b, c: (a \cdot c + b \leq c \Rightarrow a^* \cdot b \leq c) \quad \text{fof(star_induction}_1\text{, axiom)}$
 $\forall a, b, c: (c \cdot a + b \leq c \Rightarrow b \cdot a^* \leq c) \quad \text{fof(star_induction}_2\text{, axiom)}$
 $\forall a: \text{strong_iteration}(a) = a \cdot \text{strong_iteration}(a) + 1 \quad \text{fof(infty_unfold}_1\text{, axiom)}$
 $\forall a, b, c: (c \leq a \cdot c + b \Rightarrow c \leq \text{strong_iteration}(a) \cdot b) \quad \text{fof(infty_coinduction, axiom)}$
 $\forall a: \text{strong_iteration}(a) = a^* + \text{strong_iteration}(a) \cdot 0 \quad \text{fof(isolation, axiom)}$
 $\forall a, b: (a \leq b \iff a + b = b) \quad \text{fof(order, axiom)}$

KLE problems

KLE001+1.p Addition is isotone

include('Axioms/KLE001+0.ax')

$\forall x_0, x_1, x_2: (x_0 \leq x_1 \Rightarrow x_0 + x_2 \leq x_0 + x_2) \quad \text{fof(goals, conjecture)}$

KLE002+1.p Multiplication is isotone

```
include('Axioms/KLE001+0.ax')
 $\forall x_0, x_1, x_2: (x_0 \leq x_1 \Rightarrow x_0 \cdot x_2 \leq x_1 \cdot x_2)$  fof(goals, conjecture)
```

KLE003+1.p A semiring is idempotent iff 1 is idempotent

```
include('Axioms/KLE001+0.ax')
 $\forall x_0: x_0 + x_0 = x_0 \iff 1 + 1 = 1$  fof(goals, conjecture)
```

KLE004+1.p Complement - 1 is the complement of 0

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $c(0) = 1$  fof(goals, conjecture)
```

KLE005+1.p Complement - 0 is the complement of 1

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $c(1) = 0$  fof(goals, conjecture)
```

KLE006+1.p Split 1 with p

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0: (\text{test}(x_0) \Rightarrow 1 = x_0 + c(x_0))$  fof(goals, conjecture)
```

KLE007+1.p Split 1 with q and split the parts with p

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow 1 = (x_0 + c(x_0)) \cdot x_1 + (x_0 + c(x_0)) \cdot c(x_1))$  fof(goals, conjecture)
```

KLE007+2.p Split 1 with q and split the parts with p

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow (1 \leq (x_0 + c(x_0)) \cdot x_1 + (x_0 + c(x_0)) \cdot c(x_1) \text{ and } (x_0 + c(x_0)) \cdot x_1 + (x_0 + c(x_0)) \cdot c(x_1) \leq 1))$  fof(goals, conjecture)
```

KLE007+3.p Split 1 with q and split the parts with p

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow 1 = (x_0 + c(x_0)) \cdot x_1 + (x_0 + c(x_0)) \cdot c(x_1))$  fof(goals, conjecture)
```

KLE007+4.p Split 1 with q and split the parts with p

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow (1 \leq (x_0 + c(x_0)) \cdot x_1 + (x_0 + c(x_0)) \cdot c(x_1) \text{ and } (x_0 + c(x_0)) \cdot x_1 + (x_0 + c(x_0)) \cdot c(x_1) \leq 1))$  fof(goals, conjecture)
```

KLE008+1.p A simple law to eliminate the complement

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1, x_2: (\text{test}(x_2) \Rightarrow (x_0 \leq x_2 \cdot x_1 \iff (x_0 \leq x_1 \text{ and } c(x_2) \cdot x_0 \leq 0)))$  fof(goals, conjecture)
```

KLE009+1.p Split 1 into all combinations of p,q and their complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow 1 = ((x_0 \cdot x_1 + x_0 \cdot c(x_1)) + c(x_0) \cdot x_1) + c(x_0) \cdot c(x_1))$  fof(goals, conjecture)
```

KLE009+2.p Split 1 into all combinations of p,q and their complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow 1 = ((x_0 \cdot x_1 + x_0 \cdot c(x_1)) + c(x_0) \cdot x_1) + c(x_0) \cdot c(x_1))$  fof(goals, conjecture)
```

KLE009+3.p Split 1 into all combinations of p,q and their complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
```

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow (1 \leq ((x_0 \cdot x_1 + x_0 \cdot c(x_1)) + c(x_0) \cdot x_1) + c(x_0) \cdot c(x_1) \text{ and } ((x_0 \cdot x_1 + x_0 \cdot c(x_1)) + c(x_0) \cdot x_1) + c(x_0) \cdot c(x_1) \leq 1)) \quad \text{fof(goals, conjecture)}$

KLE009+4.p Split 1 into all combinations of p,q and their complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
```

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow (1 \leq ((x_0 \cdot x_1 + x_0 \cdot c(x_1)) + c(x_0) \cdot x_1) + c(x_0) \cdot c(x_1) \text{ and } ((x_0 \cdot x_1 + x_0 \cdot c(x_1)) + c(x_0) \cdot x_1) + c(x_0) \cdot c(x_1) \leq 1)) \quad \text{fof(goals, conjecture)}$

KLE010+1.p Split 1 into all combinations of p,q and their complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
```

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow 1 = (((x_1 \cdot x_0 + c(x_1) \cdot x_0) + x_0 \cdot x_1) + c(x_0) \cdot x_1) + c(x_0) \cdot c(x_1)) \quad \text{fof(goals, conjecture)}$

KLE010+2.p Split 1 into all combinations of p,q and their complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
```

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow (1 \leq (((x_1 \cdot x_0 + c(x_1) \cdot x_0) + x_0 \cdot x_1) + c(x_0) \cdot x_1) + c(x_0) \cdot c(x_1) \text{ and } (((x_1 \cdot x_0 + c(x_1) \cdot x_0) + x_0 \cdot x_1) + c(x_0) \cdot x_1) + c(x_0) \cdot c(x_1) \leq 1)) \quad \text{fof(goals, conjecture)}$

KLE010+3.p Split 1 into all combinations of p,q and their complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
```

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow 1 = (((x_1 \cdot x_0 + c(x_1) \cdot x_0) + x_0 \cdot x_1) + c(x_0) \cdot x_1) + c(x_0) \cdot c(x_1)) \quad \text{fof(goals, conjecture)}$

KLE010+4.p Split 1 into all combinations of p,q and their complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
```

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow (1 \leq (((x_1 \cdot x_0 + c(x_1) \cdot x_0) + x_0 \cdot x_1) + c(x_0) \cdot x_1) + c(x_0) \cdot c(x_1) \text{ and } (((x_1 \cdot x_0 + c(x_1) \cdot x_0) + x_0 \cdot x_1) + c(x_0) \cdot x_1) + c(x_0) \cdot c(x_1) \leq 1)) \quad \text{fof(goals, conjecture)}$

KLE011+1.p Split 1 into p,q and the product of their complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
```

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow 1 = ((x_1 + c(x_1)) \cdot x_0 + (x_0 + c(x_0)) \cdot x_1) + c(x_0) \cdot c(x_1)) \quad \text{fof(goals, conjecture)}$

KLE011+2.p Split 1 into p,q and the product of their complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
```

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow (1 \leq ((x_1 + c(x_1)) \cdot x_0 + (x_0 + c(x_0)) \cdot x_1) + c(x_0) \cdot c(x_1) \text{ and } ((x_1 + c(x_1)) \cdot x_0 + (x_0 + c(x_0)) \cdot x_1) + c(x_0) \cdot c(x_1) \leq 1)) \quad \text{fof(goals, conjecture)}$

KLE011+3.p Split 1 into p,q and the product of their complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
```

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow 1 = ((x_1 + c(x_1)) \cdot x_0 + (x_0 + c(x_0)) \cdot x_1) + c(x_0) \cdot c(x_1)) \quad \text{fof(goals, conjecture)}$

KLE011+4.p Split 1 into p,q and the product of their complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
```

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow (1 \leq ((x_1 + c(x_1)) \cdot x_0 + (x_0 + c(x_0)) \cdot x_1) + c(x_0) \cdot c(x_1) \text{ and } ((x_1 + c(x_1)) \cdot x_0 + (x_0 + c(x_0)) \cdot x_1) + c(x_0) \cdot c(x_1) \leq 1)) \quad \text{fof(goals, conjecture)}$

KLE012+1.p The multiplication of tests is commutative

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
```

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow x_0 \cdot x_1 = x_1 \cdot x_0) \quad \text{fof(goals, conjecture)}$

KLE013+1.p The complement of a product is the sum of the complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
```

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow 1 = (x_0 + x_1) + c(x_0) \cdot c(x_1)) \quad \text{fof(goals, conjecture)}$

KLE014+2.p The complement of a product is the sum of the complements

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow c(x_0 + x_1) = c(x_0) \cdot c(x_1)) \quad \text{fof(goals, conjecture)}$

KLE014+3.p The complement of a product is the sum of the complements

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow (c(x_0 + x_1) \leq c(x_0) \cdot c(x_1) \text{ and } c(x_0) \cdot c(x_1) \leq c(x_0 + x_1))) \quad \text{fof(goals, conjecture)}$

KLE015+1.p The complement of a product is the sum of the complements

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow ((x_0 + x_1) \cdot c(x_0)) \cdot c(x_1) = 0) \quad \text{fof(goals, conjecture)}$

KLE016+2.p The complement of a product is the sum of the complements

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow c(x_0 \cdot x_1) = c(x_0) + c(x_1)) \quad \text{fof(goals, conjecture)}$

KLE016+3.p The complement of a product is the sum of the complements

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1: ((\text{test}(x_1) \text{ and } \text{test}(x_0)) \Rightarrow (c(x_0 \cdot x_1) \leq c(x_0) + c(x_1) \text{ and } c(x_0) + c(x_1) \leq c(x_0 \cdot x_1))) \quad \text{fof(goals, conjecture)}$

KLE017+1.p Product of tests is their meet

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1, x_2: ((\text{test}(x_0) \text{ and } \text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow (x_2 \leq x_0 \cdot x_1 \iff (x_2 \leq x_0 \text{ and } x_2 \leq x_1))) \quad \text{fof(goals, conjecture)}$

KLE018+1.p Move a term from the left of an implication to the right and back

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1, x_2: ((\text{test}(x_0) \text{ and } \text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow (x_0 \cdot c(x_1) \leq x_2 \Rightarrow x_0 \leq x_1 + x_2)) \quad \text{fof(goals, conjecture)}$

KLE019+1.p Move a term from the left of an implication to the right and back

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1, x_2: ((\text{test}(x_0) \text{ and } \text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow (x_0 \cdot c(x_1) \leq x_2 \Leftarrow x_0 \leq x_1 + x_2)) \quad \text{fof(goals, conjecture)}$

KLE020+1.p On tests addition distributes over multiplication

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1, x_2: ((\text{test}(x_0) \text{ and } \text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow x_0 + x_1 \cdot x_2 = (x_0 + x_1) \cdot (x_0 + x_2)) \quad \text{fof(goals, conjecture)}$

KLE020+2.p On tests addition distributes over multiplication

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1, x_2: ((\text{test}(x_0) \text{ and } \text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow (x_0 + x_1 \cdot x_2 \leq (x_0 + x_1) \cdot (x_0 + x_2) \text{ and } (x_0 + x_1) \cdot (x_0 + x_2) \leq x_0 + x_1 \cdot x_2)) \quad \text{fof(goals, conjecture)}$

KLE021+1.p Decompose splitting

Decompose x into the part starting in p and the one starting in p's complement; also the rule if p then x else x = x where if p then y else z = p;y + c(p);z.

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1: (\text{test}(x_1) \Rightarrow x_0 = x_1 \cdot x_0 + c(x_1) \cdot x_0) \quad \text{fof(goals, conjecture)}$

KLE021+2.p Decompose splitting

Decompose x into the part starting in p and the one starting in p's complement; also the rule if p then x else x = x where if p then y else z = p;y + c(p);z.

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1: (\text{test}(x_1) \Rightarrow (x_0 \leq x_1 \cdot x_0 + c(x_1) \cdot x_0 \text{ and } x_1 \cdot x_0 + c(x_1) \cdot x_0 \leq x_0)) \quad \text{fof(goals, conjecture)}$

KLE021+3.p Decompose splitting

Decompose x into the part starting in p and the one starting in p 's complement; also the rule if p then x else $x = x$ where if p then y else $z = p;y + c(p);z$.

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1: (\text{test}(x_1) \Rightarrow x_0 = x_1 \cdot x_0 + c(x_1) \cdot x_0)$  fof(goals, conjecture)
```

KLE021+4.p Decompose splitting

Decompose x into the part starting in p and the one starting in p 's complement; also the rule if p then x else $x = x$ where if p then y else $z = p;y + c(p);z$.

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1: (\text{test}(x_1) \Rightarrow (x_0 \leq x_1 \cdot x_0 + c(x_1) \cdot x_0 \text{ and } x_1 \cdot x_0 + c(x_1) \cdot x_0 \leq x_0))$  fof(goals, conjecture)
```

KLE022+1.p Decompose x into parts ending in p and p 's complement

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1: (\text{test}(x_1) \Rightarrow x_0 = x_0 \cdot x_1 + x_0 \cdot c(x_1))$  fof(goals, conjecture)
```

KLE022+2.p Decompose x into parts ending in p and p 's complement

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1: (\text{test}(x_1) \Rightarrow (x_0 \leq x_0 \cdot x_1 + x_0 \cdot c(x_1) \text{ and } x_0 \cdot x_1 + x_0 \cdot c(x_1) \leq x_0))$  fof(goals, conjecture)
```

KLE022+3.p Decompose x into parts ending in p and p 's complement

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1: (\text{test}(x_1) \Rightarrow x_0 = x_0 \cdot x_1 + x_0 \cdot c(x_1))$  fof(goals, conjecture)
```

KLE022+4.p Decompose x into parts ending in p and p 's complement

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1: (\text{test}(x_1) \Rightarrow (x_0 \leq x_0 \cdot x_1 + x_0 \cdot c(x_1) \text{ and } x_0 \cdot x_1 + x_0 \cdot c(x_1) \leq x_0))$  fof(goals, conjecture)
```

KLE023+1.p Two ways of expressing the Hoare triple pxq

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1, x_2: ((\text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow (x_1 \cdot x_0 + x_0 \cdot x_2 = x_0 \cdot x_2 \Rightarrow x_0 \cdot c(x_2) + c(x_1) \cdot x_0 = c(x_1) \cdot x_0))$  fof(goals, conjecture)
```

KLE023+2.p Two ways of expressing the Hoare triple pxq

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1, x_2: ((\text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow (x_1 \cdot x_0 + x_0 \cdot x_2 = x_0 \cdot x_2 \Rightarrow x_0 \cdot c(x_2) + c(x_1) \cdot x_0 = c(x_1) \cdot x_0))$  fof(goals, conjecture)
```

KLE024+1.p Two ways of expressing the Hoare triple pxq

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1, x_2: ((\text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow (x_0 \cdot c(x_2) + c(x_1) \cdot x_0 = c(x_1) \cdot x_0 \Rightarrow (x_1 \cdot x_0) \cdot c(x_2) = 0))$  fof(goals, conjecture)
```

KLE024+2.p Two ways of expressing the Hoare triple pxq

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1, x_2: ((\text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow (x_0 \cdot c(x_2) + c(x_1) \cdot x_0 = c(x_1) \cdot x_0 \Rightarrow (x_1 \cdot x_0) \cdot c(x_2) = 0))$  fof(goals, conjecture)
```

KLE025+1.p Two ways of expressing the Hoare triple pxq

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1, x_2: ((\text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow ((x_1 \cdot x_0) \cdot c(x_2) = 0 \Rightarrow x_1 \cdot x_0 = (x_1 \cdot x_0) \cdot x_2))$  fof(goals, conjecture)
```

KLE025+2.p Two ways of expressing the Hoare triple pxq

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1, x_2: ((\text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow ((x_1 \cdot x_0) \cdot c(x_2) = 0 \Rightarrow x_1 \cdot x_0 = (x_1 \cdot x_0) \cdot x_2)) \quad \text{fof(goals, conjecture)}$ 

```

KLE026+1.p Two ways of expressing the Hoare triple pxq

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1, x_2: ((\text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow (x_1 \cdot x_0 = (x_1 \cdot x_0) \cdot x_2 \Rightarrow x_1 \cdot x_0 \leq x_0 \cdot x_2)) \quad \text{fof(goals, conjecture)}$ 

```

KLE026+2.p Two ways of expressing the Hoare triple pxq

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1, x_2: ((\text{test}(x_1) \text{ and } \text{test}(x_2)) \Rightarrow (x_1 \cdot x_0 = (x_1 \cdot x_0) \cdot x_2 \Rightarrow x_1 \cdot x_0 \leq x_0 \cdot x_2)) \quad \text{fof(goals, conjecture)}$ 

```

KLE027+1.p Simplify conditional

Simplify conditional: if p then (if p then x else y) else z = if p then x else z.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1, x_2, x_3, x_4: ((\text{test}(x_3) \text{ and } \text{test}(x_4)) \Rightarrow x_3 \cdot (x_3 \cdot x_0 + c(x_3) \cdot x_1) + c(x_3) \cdot x_2 = x_3 \cdot x_0 + c(x_3) \cdot x_2) \quad \text{fof(goals, conjecture)}$ 

```

KLE027+2.p Simplify conditional

Simplify conditional: if p then (if p then x else y) else z = if p then x else z.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1, x_2, x_3, x_4: ((\text{test}(x_3) \text{ and } \text{test}(x_4)) \Rightarrow (x_3 \cdot (x_3 \cdot x_0 + c(x_3) \cdot x_1) + c(x_3) \cdot x_2 \leq x_3 \cdot x_0 + c(x_3) \cdot x_2 \text{ and } x_3 \cdot x_0 + c(x_3) \cdot x_2 \leq x_3 \cdot (x_3 \cdot x_0 + c(x_3) \cdot x_1) + c(x_3) \cdot x_2)) \quad \text{fof(goals, conjecture)}$ 

```

KLE027+3.p Simplify conditional

Simplify conditional: if p then (if p then x else y) else z = if p then x else z.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1, x_2, x_3, x_4: ((\text{test}(x_3) \text{ and } \text{test}(x_4)) \Rightarrow x_3 \cdot (x_3 \cdot x_0 + c(x_3) \cdot x_1) + c(x_3) \cdot x_2 = x_3 \cdot x_0 + c(x_3) \cdot x_2) \quad \text{fof(goals, conjecture)}$ 

```

KLE027+4.p Simplify conditional

Simplify conditional: if p then (if p then x else y) else z = if p then x else z.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1, x_2, x_3, x_4: ((\text{test}(x_3) \text{ and } \text{test}(x_4)) \Rightarrow (x_3 \cdot (x_3 \cdot x_0 + c(x_3) \cdot x_1) + c(x_3) \cdot x_2 \leq x_3 \cdot x_0 + c(x_3) \cdot x_2 \text{ and } x_3 \cdot x_0 + c(x_3) \cdot x_2 \leq x_3 \cdot (x_3 \cdot x_0 + c(x_3) \cdot x_1) + c(x_3) \cdot x_2)) \quad \text{fof(goals, conjecture)}$ 

```

KLE028+1.p Switch nested conditions and rearrange branches of conditional

If p then (if q then u else x) else (if q then y else z) = if q then (if p then u else y) else (if p then x else z).

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1, x_2, x_3, x_4, x_5: ((\text{test}(x_4) \text{ and } \text{test}(x_5)) \Rightarrow x_4 \cdot (x_5 \cdot x_0 + c(x_5) \cdot x_1) + c(x_4) \cdot (x_5 \cdot x_2 + c(x_5) \cdot x_3) = x_5 \cdot (x_4 \cdot x_0 + c(x_4) \cdot x_2) + c(x_5) \cdot (x_4 \cdot x_1 + c(x_4) \cdot x_3)) \quad \text{fof(goals, conjecture)}$ 

```

KLE028+2.p Switch nested conditions and rearrange branches of conditional

If p then (if q then u else x) else (if q then y else z) = if q then (if p then u else y) else (if p then x else z).

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
 $\forall x_0, x_1, x_2, x_3, x_4, x_5: ((\text{test}(x_4) \text{ and } \text{test}(x_5)) \Rightarrow (x_4 \cdot (x_5 \cdot x_0 + c(x_5) \cdot x_1) + c(x_4) \cdot (x_5 \cdot x_2 + c(x_5) \cdot x_3) \leq x_5 \cdot (x_4 \cdot x_0 + c(x_4) \cdot x_2) + c(x_5) \cdot (x_4 \cdot x_1 + c(x_4) \cdot x_3) \text{ and } x_5 \cdot (x_4 \cdot x_0 + c(x_4) \cdot x_2) + c(x_5) \cdot (x_4 \cdot x_1 + c(x_4) \cdot x_3) \leq x_4 \cdot (x_5 \cdot x_0 + c(x_5) \cdot x_1) + c(x_4) \cdot (x_5 \cdot x_2 + c(x_5) \cdot x_3))) \quad \text{fof(goals, conjecture)}$ 

```

KLE028+3.p Switch nested conditions and rearrange branches of conditional

If p then (if q then u else x) else (if q then y else z) = if q then (if p then u else y) else (if p then x else z).

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')

```

$\forall x_0, x_1, x_2, x_3, x_4, x_5: ((\text{test}(x_4) \text{ and } \text{test}(x_5)) \Rightarrow x_4 \cdot (x_5 \cdot x_0 + c(x_5) \cdot x_1) + c(x_4) \cdot (x_5 \cdot x_2 + c(x_5) \cdot x_3) = x_5 \cdot (x_4 \cdot x_0 + c(x_4) \cdot x_2) + c(x_5) \cdot (x_4 \cdot x_1 + c(x_4) \cdot x_3)) \quad \text{fof(goals, conjecture)}$

KLE028+4.p Switch nested conditions and rearrange branches of conditional

If p then (if q then u else x) else (if q then y else z) = if q then (if p then u else y) else (if p then x else z).

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

include('Axioms/KLE001+2.ax')

$\forall x_0, x_1, x_2, x_3, x_4, x_5: ((\text{test}(x_4) \text{ and } \text{test}(x_5)) \Rightarrow (x_4 \cdot (x_5 \cdot x_0 + c(x_5) \cdot x_1) + c(x_4) \cdot (x_5 \cdot x_2 + c(x_5) \cdot x_3) \leq x_5 \cdot (x_4 \cdot x_0 + c(x_4) \cdot x_2) + c(x_5) \cdot (x_4 \cdot x_1 + c(x_4) \cdot x_3) \text{ and } x_5 \cdot (x_4 \cdot x_0 + c(x_4) \cdot x_2) + c(x_5) \cdot (x_4 \cdot x_1 + c(x_4) \cdot x_3) \leq x_4 \cdot (x_5 \cdot x_0 + c(x_5) \cdot x_1) + c(x_4) \cdot (x_5 \cdot x_2 + c(x_5) \cdot x_3))) \quad \text{fof(goals, conjecture)}$

KLE029+1.p Characterisations of meet

Equivalence of characterisation as greatest lower bound and universal characterisation.

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+3.ax')

$\forall x_0, x_1, x_2: (\text{ismeet}(x_0, x_1, x_2) \iff \text{ismeetu}(x_0, x_1, x_2)) \quad \text{fof(goals, conjecture)}$

KLE030+1.p Restriction can be pulled out of meet

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

include('Axioms/KLE001+3.ax')

$\forall x_0, x_1, x_2, x_3: ((\text{test}(x_3) \text{ and } \text{ismeet}(x_0, x_1, x_2)) \Rightarrow \text{ismeet}(x_3 \cdot x_0, x_3 \cdot x_1, x_2)) \quad \text{fof(goals, conjecture)}$

KLE031+1.p Restriction distributes through meet

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

include('Axioms/KLE001+3.ax')

$\forall x_0, x_1, x_2, x_3: ((\text{test}(x_3) \text{ and } \text{ismeet}(x_0, x_1, x_2)) \Rightarrow \text{ismeet}(x_3 \cdot x_0, x_3 \cdot x_1, x_3 \cdot x_2)) \quad \text{fof(goals, conjecture)}$

KLE032+1.p Meet of restrictions is restriction by meet

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

include('Axioms/KLE001+3.ax')

$\forall x_0, x_1, x_2: ((\text{test}(x_2) \text{ and } \text{test}(x_1)) \Rightarrow \text{ismeet}((x_1 \cdot x_2) \cdot x_0, x_1 \cdot x_0, x_2 \cdot x_0)) \quad \text{fof(goals, conjecture)}$

KLE033+1.p Disjoint tests induce disjoint restrictions

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

include('Axioms/KLE001+3.ax')

$\forall x_0, x_1, x_2: ((\text{test}(x_2) \text{ and } \text{test}(x_1) \text{ and } \text{ismeet}(0, x_1, x_2)) \Rightarrow \text{ismeet}(0, x_1 \cdot x_0, x_2 \cdot x_0)) \quad \text{fof(goals, conjecture)}$

KLE034+1.p Hoare rule product

Encoding of Hoare rule pxq <description> qyr → px;yr.

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

include('Axioms/KLE001+2.ax')

$\forall x_0, x_1, x_2, x_3, x_4: ((\text{test}(x_3) \text{ and } \text{test}(x_2) \text{ and } \text{test}(x_4) \text{ and } (x_2 \cdot x_0) \cdot c(x_3) \leq 0 \text{ and } (x_3 \cdot x_1) \cdot c(x_4) \leq 0) \Rightarrow ((x_2 \cdot x_0) \cdot x_1) \cdot c(x_4) \leq 0) \quad \text{fof(goals, conjecture)}$

KLE034+2.p Hoare rule product

Encoding of Hoare rule pxq <description> qyr → px;yr.

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

include('Axioms/KLE001+2.ax')

$\forall x_0, x_1, x_2, x_3, x_4: ((\text{test}(x_3) \text{ and } \text{test}(x_2) \text{ and } \text{test}(x_4) \text{ and } (x_2 \cdot x_0) \cdot c(x_3) \leq 0 \text{ and } (x_3 \cdot x_1) \cdot c(x_4) \leq 0) \Rightarrow ((x_2 \cdot x_0) \cdot x_1) \cdot c(x_4) \leq 0) \quad \text{fof(goals, conjecture)}$

KLE035+1.p Hoare rule sum

Encoding of Hoare rule pxq <description> qyr → px;yr.

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+1.ax')

$\forall x_0, x_1, x_2, x_3: ((\text{test}(x_3) \text{ and } \text{test}(x_2) \text{ and } (x_2 \cdot x_0) \cdot c(x_3) \leq 0 \text{ and } (x_2 \cdot x_1) \cdot c(x_3) \leq 0) \Rightarrow (x_2 \cdot (x_0 + x_1)) \cdot c(x_3) \leq 0) \quad \text{fof(goals, conjecture)}$

KLE035+2.p Hoare rule sum

Encoding of Hoare rule pxq <description> qyr → px;yr.
 include('Axioms/KLE001+0.ax')
 include('Axioms/KLE001+1.ax')
 include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1, x_2, x_3: ((\text{test}(x_3) \text{ and } \text{test}(x_2) \text{ and } (x_2 \cdot x_0) \cdot c(x_3) \leq 0 \text{ and } (x_2 \cdot x_1) \cdot c(x_3) \leq 0) \Rightarrow (x_2 \cdot (x_0 + x_1)) \cdot c(x_3) \leq 0)$ fof(goals, conjecture)

KLE036+1.p Star recursion

Together with star unfold this yields the classical recursion equation for star.

include('Axioms/KLE002+0.ax')
 $\forall x_0: x_0^* \leq 1 + x_0 \cdot x_0^*$ fof(goals, conjecture)

KLE037+1.p Star reflexivity

Star is reflexive, i.e., contains the identity.

include('Axioms/KLE002+0.ax')
 $\forall x_0: 1 \leq x_0^*$ fof(goals, conjecture)

KLE038+1.p Star extensivity

Star is extensive.

include('Axioms/KLE002+0.ax')
 $\forall x_0: x_0 \leq x_0^*$ fof(goals, conjecture)

KLE039+1.p Star idempotence

The star operation is idempotent.

include('Axioms/KLE002+0.ax')
 $\forall x_0: (x_0^*)^* = x_0^*$ fof(goals, conjecture)

KLE039+2.p Star idempotence

The star operation is idempotent.

include('Axioms/KLE002+0.ax')
 $\forall x_0: ((x_0^*)^* \leq x_0^* \text{ and } x_0^* \leq (x_0^*)^*)$ fof(goals, conjecture)

KLE040+1.p The star of an element is multiplicatively idempotent

include('Axioms/KLE002+0.ax')
 $\forall x_0: x_0^* \cdot x_0^* = x_0^*$ fof(goals, conjecture)

KLE040+2.p The star of an element is multiplicatively idempotent

include('Axioms/KLE002+0.ax')
 $\forall x_0: (x_0^* \cdot x_0^* \leq x_0^* \text{ and } x_0^* \leq x_0^* \cdot x_0^*)$ fof(goals, conjecture)

KLE041+1.p Star isotony

The star operation is isotone.

include('Axioms/KLE002+0.ax')
 $\forall x_0, x_1: (x_0 \leq x_1 \Rightarrow x_0^* \leq x_1^*)$ fof(goals, conjecture)

KLE042+1.p Star sliding

Two ways of grouping an alternation of x's and y's that starts and ends with x.

include('Axioms/KLE002+0.ax')
 $\forall x_0, x_1: (x_0 \cdot x_1)^* \cdot x_0 = x_0 \cdot (x_1 \cdot x_0)^*$ fof(goals, conjecture)

KLE042+2.p Star sliding

Two ways of grouping an alternation of x's and y's that starts and ends with x.

include('Axioms/KLE002+0.ax')
 $\forall x_0, x_1: ((x_0 \cdot x_1)^* \cdot x_0 \leq x_0 \cdot (x_1 \cdot x_0)^* \text{ and } x_0 \cdot (x_1 \cdot x_0)^* \leq (x_0 \cdot x_1)^* \cdot x_0)$ fof(goals, conjecture)

KLE043+1.p Star recursion

Classical recursion equation for star of x followed by y.

include('Axioms/KLE002+0.ax')
 $\forall x_0, x_1: x_0^* \cdot x_1 = x_1 + (x_0 \cdot x_0^*) \cdot x_1$ fof(goals, conjecture)

KLE043+2.p Star recursion

Classical recursion equation for star of x followed by y.

include('Axioms/KLE002+0.ax')
 $\forall x_0, x_1: (x_0^* \cdot x_1 \leq x_1 + (x_0 \cdot x_0^*) \cdot x_1 \text{ and } x_1 + (x_0 \cdot x_0^*) \cdot x_1 \leq x_0^* \cdot x_1)$ fof(goals, conjecture)

KLE044+1.p Star simplification

Identity can be eliminated from the star of a sum.

```
include('Axioms/KLE002+0.ax')
 $\forall x_0: (1 + x_0)^* = x_0^*$  fof(goals, conjecture)
```

KLE044+2.p Star simplification

Identity can be eliminated from the star of a sum.

```
include('Axioms/KLE002+0.ax')
 $\forall x_0: ((1 + x_0)^* \leq x_0^* \text{ and } x_0^* \leq (1 + x_0)^*)$  fof(goals, conjecture)
```

KLE045+1.p Semi-commutation

If x semi-commutes over y transforming into z then the same applies to x and y^* .

```
include('Axioms/KLE002+0.ax')
 $\forall x_0, x_1, x_2: (x_0 \cdot x_2 \leq x_2 \cdot x_1 \Rightarrow x_0^* \cdot x_2 \leq x_2 \cdot x_1^*)$  fof(goals, conjecture)
```

KLE046+1.p Church-Rosser theorem

Normalise arbitrary sequences of x 's and y 's into x 's followed by y 's.

```
include('Axioms/KLE002+0.ax')
```

```
 $\forall x_0, x_1: (x_1^* \cdot x_0^* \leq x_0^* \cdot x_1^* \Rightarrow (x_0 + x_1)^* \leq x_0^* \cdot x_1^*)$  fof(goals, conjecture)
```

KLE047+1.p Star simplification

Semi-commutation of x and y allows normalizing arbitrary sequences of x 's and y 's.

```
include('Axioms/KLE002+0.ax')
 $\forall x_0, x_1: (x_1 \cdot x_0 \leq x_0 \cdot x_1 \Rightarrow (x_0 + x_1)^* \leq x_0^* \cdot x_1^*)$  fof(goals, conjecture)
```

KLE048+1.p The star of a test is always 1

```
include('Axioms/KLE002+0.ax')
```

```
include('Axioms/KLE001+1.ax')
```

```
 $\forall x_0: (\text{test}(x_0) \Rightarrow x_0^* = 1)$  fof(goals, conjecture)
```

KLE049+1.p Hoare rule while

Encoding of Hoare rule $p;qxq \rightarrow q$ while p do x $c(p);q$, where while p do $x = (p;x)^*;c(p)$.

```
include('Axioms/KLE002+0.ax')
```

```
include('Axioms/KLE001+1.ax')
```

```
 $\forall x_0, x_1, x_2: ((\text{test}(x_1) \text{ and } \text{test}(x_2)) \text{ and } ((x_2 \cdot x_1) \cdot x_0) \cdot c(x_2) \leq 0 \Rightarrow ((x_2 \cdot (x_1 \cdot x_0)^*) \cdot c(x_1)) \cdot c(x_2) \leq 0)$  fof(goals, conjecture)
```

KLE049+2.p Hoare rule while

Encoding of Hoare rule $p;qxq \rightarrow q$ while p do x $c(p);q$, where while p do $x = (p;x)^*;c(p)$.

```
include('Axioms/KLE002+0.ax')
```

```
include('Axioms/KLE001+1.ax')
```

```
include('Axioms/KLE001+2.ax')
```

```
 $\forall x_0, x_1, x_2: ((\text{test}(x_1) \text{ and } \text{test}(x_2)) \text{ and } ((x_2 \cdot x_1) \cdot x_0) \cdot c(x_2) \leq 0 \Rightarrow ((x_2 \cdot (x_1 \cdot x_0)^*) \cdot c(x_1)) \cdot c(x_2) \leq 0)$  fof(goals, conjecture)
```

KLE050+1.p Loop denesting

Encoding of the law while p do $(x ; \text{while } q \text{ do } y) = \text{if } p \text{ then } (x ; \text{while } p+q \text{ do if } q \text{ then } y \text{ else } x) \text{ else skip}$.

```
include('Axioms/KLE002+0.ax')
```

```
include('Axioms/KLE001+1.ax')
```

```
 $\forall x_0, x_1, x_2, x_3: ((\text{test}(x_2) \text{ and } \text{test}(x_3)) \Rightarrow (((x_2 \cdot x_0) \cdot (x_3 \cdot x_1)^*) \cdot c(x_3))^* \cdot c(x_2) = (((x_2 \cdot x_0) \cdot (x_2 + x_3)) \cdot (x_3 \cdot x_1 + c(x_3) \cdot x_0)^*) \cdot c(x_2 + x_3) + c(x_2))$  fof(goals, conjecture)
```

KLE050+2.p Loop denesting

Encoding of the law while p do $(x ; \text{while } q \text{ do } y) = \text{if } p \text{ then } (x ; \text{while } p+q \text{ do if } q \text{ then } y \text{ else } x) \text{ else skip}$.

```
include('Axioms/KLE002+0.ax')
```

```
include('Axioms/KLE001+1.ax')
```

```
 $\forall x_0, x_1, x_2, x_3: ((\text{test}(x_2) \text{ and } \text{test}(x_3)) \Rightarrow ((((x_2 \cdot x_0) \cdot (x_3 \cdot x_1)^*) \cdot c(x_3))^* \cdot c(x_2) \leq (((x_2 \cdot x_0) \cdot (x_2 + x_3)) \cdot (x_3 \cdot x_1 + c(x_3) \cdot x_0)^*) \cdot c(x_2 + x_3) + c(x_2) \leq (((x_2 \cdot x_0) \cdot (x_3 \cdot x_1)^*) \cdot c(x_3))^* \cdot c(x_2))$  fof(goals, conjecture)
```

KLE050+3.p Loop denesting

Encoding of the law while p do $(x ; \text{while } q \text{ do } y) = \text{if } p \text{ then } (x ; \text{while } p+q \text{ do if } q \text{ then } y \text{ else } x) \text{ else skip}$.

```
include('Axioms/KLE002+0.ax')
```

```
include('Axioms/KLE001+1.ax')
```

```
include('Axioms/KLE001+2.ax')
```

```
 $\forall x_0, x_1, x_2, x_3: ((\text{test}(x_2) \text{ and } \text{test}(x_3)) \Rightarrow (((x_2 \cdot x_0) \cdot (x_3 \cdot x_1)^*) \cdot c(x_3))^* \cdot c(x_2) = (((x_2 \cdot x_0) \cdot (x_2 + x_3)) \cdot (x_3 \cdot x_1 + c(x_3) \cdot x_0)^*) \cdot c(x_2 + x_3) + c(x_2))$  fof(goals, conjecture)
```

KLE050+4.p Loop denesting

Encoding of the law while p do $(x ; \text{while } q \text{ do } y) = \text{if } p \text{ then } (x ; \text{while } p+q \text{ do if } q \text{ then } y \text{ else } x) \text{ else skip}$.

```

include('Axioms/KLE002+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
 $\forall x_0, x_1, x_2, x_3: ((\text{test}(x_2) \text{ and } \text{test}(x_3)) \Rightarrow (((x_2 \cdot x_0) \cdot (x_3 \cdot x_1)^*) \cdot c(x_3))^* \cdot c(x_2) \leq (((x_2 \cdot x_0) \cdot (x_2 + x_3)) \cdot (x_3 \cdot x_1 + c(x_3) \cdot x_0)^*) \cdot c(x_2 + x_3) + c(x_2) \leq (((x_2 \cdot x_0) \cdot (x_3 \cdot x_1)^*) \cdot c(x_3))^* \cdot c(x_2)) \quad \text{fof(goals, conjecture)}$ 

```

KLE051+1.p Every domain semiring is automatically isotone

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0: x_0 + x_0 = x_0 \quad \text{fof(goals, conjecture)}$ 

```

KLE052+1.p Domain is a left invariant

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0: \text{dom}(x_0) \cdot x_0 = x_0 \quad \text{fof(goals, conjecture)}$ 

```

KLE053+1.p Domain is a projection

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0: \text{dom}(\text{dom}(x_0)) = \text{dom}(x_0) \quad \text{fof(goals, conjecture)}$ 

```

KLE054+1.p Domain is prefix increasing

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: \text{dom}(x_0 \cdot x_1) + \text{dom}(x_0) = \text{dom}(x_0) \quad \text{fof(goals, conjecture)}$ 

```

KLE055+1.p Domain expands subidentities

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0: (x_0 + 1 = 1 \Rightarrow x_0 + \text{dom}(x_0) = \text{dom}(x_0)) \quad \text{fof(goals, conjecture)}$ 

```

KLE056+1.p Domain is very strict

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0: (\text{dom}(x_0) = 0 \Rightarrow x_0 = 0) \quad \text{fof(goals, conjecture)}$ 

```

KLE057+1.p Domain is very strict

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0: (\text{dom}(x_0) = 0 \Leftarrow x_0 = 0) \quad \text{fof(goals, conjecture)}$ 

```

KLE058+1.p Domain is costrict

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\text{dom}(1) = 1 \quad \text{fof(goals, conjecture)}$ 

```

KLE059+1.p Domain is isotone

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: (x_0 + x_1 = x_1 \Rightarrow \text{dom}(x_0) + \text{dom}(x_1) = \text{dom}(x_1)) \quad \text{fof(goals, conjecture)}$ 

```

KLE060+1.p Domain elements can be exported

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: \text{dom}(\text{dom}(x_0) \cdot x_1) = \text{dom}(x_0) \cdot \text{dom}(x_1) \quad \text{fof(goals, conjecture)}$ 

```

KLE061+1.p Domain elements are multiplicatively idempotent

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0: \text{dom}(x_0) \cdot \text{dom}(x_0) = \text{dom}(x_0) \quad \text{fof(goals, conjecture)}$ 

```

KLE062+1.p Domain elements are multiplicatively commutative

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: \text{dom}(x_0) \cdot \text{dom}(x_1) = \text{dom}(x_1) \cdot \text{dom}(x_0) \quad \text{fof(goals, conjecture)}$ 

```

KLE063+1.p Domain elements are least left preservers

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: (x_0 + \text{dom}(x_1) \cdot x_0 = \text{dom}(x_1) \cdot x_0 \Rightarrow \text{dom}(x_0) + \text{dom}(x_1) = \text{dom}(x_1))$  fof(goals, conjecture)
```

KLE064+1.p Domain elements are least left preservers

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: (x_0 + \text{dom}(x_1) \cdot x_0 = \text{dom}(x_1) \cdot x_0 \Leftarrow \text{dom}(x_0) + \text{dom}(x_1) = \text{dom}(x_1))$  fof(goals, conjecture)
```

KLE065+1.p Domain is weakly local

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: (x_0 \cdot x_1 = 0 \Rightarrow x_0 \cdot \text{dom}(x_1) = 0)$  fof(goals, conjecture)
```

KLE066+1.p Domain is weakly local

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: (x_0 \cdot x_1 = 0 \Leftarrow x_0 \cdot \text{dom}(x_1) = 0)$  fof(goals, conjecture)
```

KLE067+1.p Domain elements are closed under addition

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: \text{dom}(\text{dom}(x_0) + \text{dom}(x_1)) = \text{dom}(x_0) + \text{dom}(x_1)$  fof(goals, conjecture)
```

KLE068+1.p Domain elements are closed under multiplication

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: \text{dom}(\text{dom}(x_0) \cdot \text{dom}(x_1)) = \text{dom}(x_0) \cdot \text{dom}(x_1)$  fof(goals, conjecture)
```

KLE069+1.p Domain elements satisfy the first lattice absorption law

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: \text{dom}(x_0) \cdot (\text{dom}(x_0) + \text{dom}(x_1)) = \text{dom}(x_0)$  fof(goals, conjecture)
```

KLE070+1.p Domain elements satisfy the second lattice absorption law

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: \text{dom}(x_0) + \text{dom}(x_0) \cdot \text{dom}(x_1) = \text{dom}(x_0)$  fof(goals, conjecture)
```

KLE071+1.p Domain elements satisfy one of the lattice distributivity laws

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1, x_2: \text{dom}(x_0) + \text{dom}(x_1) \cdot \text{dom}(x_2) = (\text{dom}(x_0) + \text{dom}(x_1)) \cdot (\text{dom}(x_0) + \text{dom}(x_2))$  fof(goals, conjecture)
```

KLE072+1.p Domain elements satisfy the first axiom of Kleene modules

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1, x_2: \text{dom}((x_0 + x_1) \cdot \text{dom}(x_2)) = \text{dom}(x_0 \cdot \text{dom}(x_2)) + \text{dom}(x_1 \cdot \text{dom}(x_2))$  fof(goals, conjecture)
```

KLE073+1.p Domain elements satisfy the second Kleene module axiom

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
 $\forall x_0, x_1, x_2: \text{dom}(x_0 \cdot (\text{dom}(x_1) + \text{dom}(x_2))) = \text{dom}(x_0 \cdot \text{dom}(x_1)) + \text{dom}(x_0 \cdot \text{dom}(x_2))$  fof(goals, conjecture)
```

KLE074+1.p Domain elements satisfy the third Kleene module axiom.

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1, x_2: \text{dom}((x_0 \cdot x_1) \cdot \text{dom}(x_2)) = \text{dom}(x_0 \cdot \text{dom}(x_1 \cdot \text{dom}(x_2)))$  fof(goals, conjecture)
```

KLE075+1.p Domain elements satisfy the fourth Kleene module axiom

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0: \text{dom}(1 \cdot \text{dom}(x_0)) = \text{dom}(x_0)$  fof(goals, conjecture)
```

KLE076+1.p Domain elements satisfy the fifth Kleene module axiom

```
include('Axioms/KLE001+0.ax')
```

```
include('Axioms/KLE001+5.ax')
 $\forall x_0: \text{dom}(x_0 \cdot \text{dom}(0)) = 0$  fof(goals, conjecture)
```

KLE077+1.p Domain elements satisfy the sixth Kleene module axiom

```
include('Axioms/KLE002+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1, x_2: ((\text{dom}(x_0) + \text{dom}(x_1 \cdot \text{dom}(x_2))) + \text{dom}(x_2)) = \text{dom}(x_2) \Rightarrow \text{dom}(x_1^* \cdot \text{dom}(x_0)) + \text{dom}(x_2) = \text{dom}(x_2)$  fof(goals, conjecture)
```

KLE078+1.p Antidomain elements are domain elements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0: (\forall x_1: (\text{dom}(x_1) + \text{ad}(x_1) = 1 \text{ and } \text{dom}(x_1) \cdot \text{ad}(x_1) = 0) \Rightarrow \text{dom}(\text{ad}(x_0)) = \text{ad}(x_0))$  fof(goals, conjecture)
```

KLE079+1.p Domain and antidomain elements are complements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0: (\forall x_1: (\text{dom}(x_1) + \text{ad}(x_1) = 1 \text{ and } \text{dom}(x_1) \cdot \text{ad}(x_1) = 0) \Rightarrow \text{ad}(x_0) \cdot \text{dom}(x_0) = 0)$  fof(goals, conjecture)
```

KLE080+1.p Another complementation property of domain and antidomain elements

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0: (\forall x_1: (\text{dom}(x_1) + \text{ad}(x_1) = 1 \text{ and } \text{dom}(x_1) \cdot \text{ad}(x_1) = 0) \Rightarrow \text{ad}(\text{ad}(x_0)) = \text{dom}(x_0))$  fof(goals, conjecture)
```

KLE081+1.p Antidomain elements are left annihilators

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0: (\forall x_1: (\text{dom}(x_1) + \text{ad}(x_1) = 1 \text{ and } \text{dom}(x_1) \cdot \text{ad}(x_1) = 0) \Rightarrow \text{ad}(x_0) \cdot x_0 = 0)$  fof(goals, conjecture)
```

KLE082+1.p Antidomain is local with respect to multiplication

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
 $\forall x_0, x_1: (\forall x_2: (\text{dom}(x_2) + \text{ad}(x_2) = 1 \text{ and } \text{dom}(x_2) \cdot \text{ad}(x_2) = 0) \Rightarrow \text{ad}(x_0 \cdot x_1) + \text{ad}(x_0 \cdot \text{dom}(x_1)) = \text{ad}(x_0 \cdot \text{dom}(x_1)))$  fof(goals, conjecture)
```

KLE083+1.p Domain is a left invariant

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
 $\forall x_0: x_0 = \text{dom}(x_0) \cdot x_0$  fof(goals, conjecture)
```

KLE084+1.p Domain is local with respect to multiplication

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
 $\forall x_0, x_1: \text{dom}(x_0 \cdot x_1) = \text{dom}(x_0 \cdot \text{dom}(x_1))$  fof(goals, conjecture)
```

KLE085+1.p Domain elements are subidenties

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
 $\forall x_0: \text{dom}(x_0) + 1 = 1$  fof(goals, conjecture)
```

KLE086+1.p Domain is strict

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
 $\text{dom}(0) = 0$  fof(goals, conjecture)
```

KLE087+1.p Domain is additive

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
 $\forall x_0, x_1: \text{dom}(x_0 + x_1) = \text{dom}(x_0) + \text{dom}(x_1)$  fof(goals, conjecture)
```

KLE088+1.p Antidomain elements are greatest left annihilators

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
 $\forall x_0, x_1: (\text{dom}(x_0) \cdot x_1 = 0 \Rightarrow \text{dom}(x_0) + \text{ad}(x_1) = \text{ad}(x_1))$  fof(goals, conjecture)
```

KLE089+1.p Antidomain elements are greatest left annihilators

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
```

$\forall x_0, x_1: (\text{dom}(x_0) \cdot x_1 = 0 \Leftarrow \text{dom}(x_0) + \text{ad}(x_1) = \text{ad}(x_1)) \quad \text{fof(goals, conjecture)}$

KLE090+1.p Antidomain elements are antitone

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

$\forall x_0, x_1: (x_0 + x_1 = x_1 \Rightarrow \text{ad}(x_1) + \text{ad}(x_0) = \text{ad}(x_0)) \quad \text{fof(goals, conjecture)}$

KLE091+1.p Codomain closure

In a Boolean domain semiring, codomain elements are domain elements.

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

$\forall x_0: \text{dom}(\text{cod}(x_0)) = \text{cod}(x_0) \quad \text{fof(goals, conjecture)}$

KLE092+1.p Coantidomain closure

In a Boolean domain semiring, coantidomain elements are domain elements.

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

$\forall x_0: \text{dom}(\text{coad}(x_0)) = \text{coad}(x_0) \quad \text{fof(goals, conjecture)}$

KLE093+1.p Domain of star

In a Kleene algebra with domain, the domain of a star is always one.

include('Axioms/KLE002+0.ax')

include('Axioms/KLE001+5.ax')

$\forall x_0: \text{dom}(x_0^*) = 1 \quad \text{fof(goals, conjecture)}$

KLE094+1.p Segerberg

Segerberg's formula holds in every Kleene algebra with Boolean domain.

include('Axioms/KLE002+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

$\forall x_0, x_1: \text{domain_difference}(\text{forward_diamond}(x_0^*, \text{dom}(x_1)), \text{dom}(x_1)) + \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{forward_diamond}(x_0, \text{dom}(x_1)), \text{dom}(x_1))) \quad \text{fof(goals, conjecture)}$

KLE095+1.p Modal operators satisfy a star unfold law

include('Axioms/KLE002+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

$\forall x_0, x_1: \text{dom}(x_0) + \text{forward_diamond}(x_1, \text{forward_diamond}(x_0^*, \text{dom}(x_0))) = \text{forward_diamond}(x_1^*, \text{dom}(x_0)) \quad \text{fof(goals, conjecture)}$

KLE096+1.p Modal operators satisfy a star unfold law

include('Axioms/KLE002+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

$\forall x_0, x_1: \text{dom}(x_0) + \text{forward_diamond}(x_1^*, \text{forward_diamond}(x_1, \text{dom}(x_0))) = \text{forward_diamond}(x_1^*, \text{dom}(x_0)) \quad \text{fof(goals, conjecture)}$

KLE097+1.p Modal operators at left hand-sides can be eliminated

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

$\forall x_0, x_1, x_2: (\text{forward_diamond}(x_0, \text{dom}(x_1)) + \text{dom}(x_2) = \text{dom}(x_2) \Leftarrow x_0 \cdot \text{dom}(x_1) + \text{dom}(x_2) \cdot x_0 = \text{dom}(x_2) \cdot x_0) \quad \text{fof(goals, conjecture)}$

KLE098+1.p Modal operators at left hand-sides can be eliminated

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

$\forall x_0, x_1, x_2: (\text{forward_diamond}(x_0, \text{dom}(x_1)) + \text{dom}(x_2) = \text{dom}(x_2) \Rightarrow x_0 \cdot \text{dom}(x_1) + \text{dom}(x_2) \cdot x_0 = \text{dom}(x_2) \cdot x_0) \quad \text{fof(goals, conjecture)}$

KLE099+1.p Modal operators at left hand-sides can be eliminated

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

$\forall x_0, x_1, x_2: (\text{forward_diamond}(x_0, \text{dom}(x_1)) + \text{dom}(x_2) = \text{dom}(x_2) \Rightarrow \text{ad}(x_2) \cdot (x_0 \cdot \text{dom}(x_1)) = 0) \quad \text{fof(goals, conjecture)}$

KLE100+1.p Modal operators at left hand-sides can be eliminated

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1, x_2: (\text{forward\_diamond}(x_0, \text{dom}(x_1)) + \text{dom}(x_2) = \text{dom}(x_2) \Leftarrow \text{ad}(x_2) \cdot (x_0 \cdot \text{dom}(x_1)) = 0)$  fof(goals, conjecture)

```

KLE101+1.p Forward and backward diamonds are conjugates

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1, x_2: (\text{forward\_diamond}(x_0, \text{dom}(x_1)) \cdot \text{dom}(x_2) = 0 \Rightarrow \text{dom}(x_1) \cdot \text{backward\_diamond}(x_0, \text{dom}(x_2)) = 0)$  fof(goals, conjecture)

```

KLE102+1.p Forward and backward diamonds are conjugates

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1, x_2: (\text{forward\_diamond}(x_0, \text{dom}(x_1)) \cdot \text{dom}(x_2) = 0 \Leftarrow \text{dom}(x_1) \cdot \text{backward\_diamond}(x_0, \text{dom}(x_2)) = 0)$  fof(goals, conjecture)

```

KLE103+1.p Forward and backward boxes are conjugates

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1, x_2: (\text{forward\_box}(x_0, \text{dom}(x_1)) + \text{dom}(x_2) = 1 \Rightarrow \text{dom}(x_1) + \text{backward\_box}(x_0, \text{dom}(x_2)) = 1)$  fof(goals, conjecture)

```

KLE104+1.p Forward and backward boxes are conjugates

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1, x_2: (\text{forward\_box}(x_0, \text{dom}(x_1)) + \text{dom}(x_2) = 1 \Leftarrow \text{dom}(x_1) + \text{backward\_box}(x_0, \text{dom}(x_2)) = 1)$  fof(goals, conjecture)

```

KLE105+1.p Galois

Forward diamonds and backward boxes are adjoints in a Galois connection.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1, x_2: (\text{forward\_diamond}(x_0, \text{dom}(x_1)) + \text{dom}(x_2) = \text{dom}(x_2) \Rightarrow \text{dom}(x_1) + \text{backward\_box}(x_0, \text{dom}(x_2)) = \text{backward\_box}(x_0, \text{dom}(x_2)))$  fof(goals, conjecture)

```

KLE106+1.p Galois

Forward diamonds and backward boxes are adjoints in a Galois connection.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1, x_2: (\text{forward\_diamond}(x_0, \text{dom}(x_1)) + \text{dom}(x_2) = \text{dom}(x_2) \Leftarrow \text{dom}(x_1) + \text{backward\_box}(x_0, \text{dom}(x_2)) = \text{backward\_box}(x_0, \text{dom}(x_2)))$  fof(goals, conjecture)

```

KLE107+1.p Galois

Backward diamonds and forward boxes are adjoints of a Galois connection.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1, x_2: (\text{backward\_diamond}(x_0, \text{dom}(x_1)) + \text{dom}(x_2) = \text{dom}(x_2) \Rightarrow \text{dom}(x_1) + \text{forward\_box}(x_0, \text{dom}(x_2)) = \text{forward\_box}(x_0, \text{dom}(x_2)))$  fof(goals, conjecture)

```

KLE108+1.p Galois

Backward diamonds and forward boxes are adjoints of a Galois connection.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1, x_2: (\text{backward\_diamond}(x_0, \text{dom}(x_1)) + \text{dom}(x_2) = \text{dom}(x_2) \Leftarrow \text{dom}(x_1) + \text{forward\_box}(x_0, \text{dom}(x_2)) = \text{forward\_box}(x_0, \text{dom}(x_2)))$  fof(goals, conjecture)

```

KLE109+1.p Forward diamonds and backward boxes satisfy a cancellation law

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')

```

$\forall x_0, x_1: \text{forward_diamond}(x_0, \text{backward_box}(x_0, \text{dom}(x_1))) + \text{dom}(x_1) = \text{dom}(x_1)$	fof(goals, conjecture)
KLE110+1.p Forward diamonds and backward boxes satisfy a cancellation law	
include('Axioms/KLE001+0.ax')	
include('Axioms/KLE001+4.ax')	
include('Axioms/KLE001+6.ax')	
$\forall x_0, x_1: \text{dom}(x_0) + \text{backward_box}(x_1, \text{forward_diamond}(x_1, \text{dom}(x_0))) = \text{backward_box}(x_1, \text{forward_diamond}(x_1, \text{dom}(x_0)))$	
KLE111+1.p Backward diamonds and forward boxes satisfy a cancellation law	
include('Axioms/KLE001+0.ax')	
include('Axioms/KLE001+4.ax')	
include('Axioms/KLE001+6.ax')	
$\forall x_0, x_1: \text{backward_diamond}(x_0, \text{forward_box}(x_0, \text{dom}(x_1))) + \text{dom}(x_1) = \text{dom}(x_1)$	fof(goals, conjecture)
KLE112+1.p Backward diamonds and forward boxes satisfy a cancellation law	
include('Axioms/KLE001+0.ax')	
include('Axioms/KLE001+4.ax')	
include('Axioms/KLE001+6.ax')	
$\forall x_0, x_1: \text{dom}(x_0) + \text{forward_box}(x_1, \text{backward_diamond}(x_1, \text{dom}(x_0))) = \text{forward_box}(x_1, \text{backward_diamond}(x_1, \text{dom}(x_0)))$	
KLE113+1.p Diamonds are strict	
include('Axioms/KLE001+0.ax')	
include('Axioms/KLE001+4.ax')	
include('Axioms/KLE001+6.ax')	
$\forall x_0: \text{forward_diamond}(x_0, 0) = 0$	fof(goals, conjecture)
KLE114+1.p Boxes are costrict	
include('Axioms/KLE001+0.ax')	
include('Axioms/KLE001+4.ax')	
include('Axioms/KLE001+6.ax')	
$\forall x_0: \text{forward_box}(x_0, 1) = 1$	fof(goals, conjecture)
KLE115+1.p Diamonds are additive	
include('Axioms/KLE001+0.ax')	
include('Axioms/KLE001+4.ax')	
include('Axioms/KLE001+6.ax')	
$\forall x_0, x_1, x_2: \text{forward_diamond}(x_0, \text{dom}(x_1) + \text{dom}(x_2)) = \text{forward_diamond}(x_0, \text{dom}(x_1)) + \text{forward_diamond}(x_0, \text{dom}(x_2))$	
KLE116+1.p Boxes are multiplicative	
include('Axioms/KLE001+0.ax')	
include('Axioms/KLE001+4.ax')	
include('Axioms/KLE001+6.ax')	
$\forall x_0, x_1, x_2: \text{forward_box}(x_0, \text{dom}(x_1) \cdot \text{dom}(x_2)) = \text{forward_box}(x_0, \text{dom}(x_1)) \cdot \text{forward_box}(x_0, \text{dom}(x_2))$	fof(goals, conjecture)
KLE117+1.p Diamonds are isotone	
include('Axioms/KLE001+0.ax')	
include('Axioms/KLE001+4.ax')	
include('Axioms/KLE001+6.ax')	
$\forall x_0, x_1: (x_0 + x_1 = x_1 \Rightarrow \forall x_2: \text{forward_diamond}(x_0, \text{dom}(x_2)) + \text{forward_diamond}(x_1, \text{dom}(x_2)) = \text{forward_diamond}(x_1, \text{dom}(x_2)))$	
KLE118+1.p Boxes are antitone	
include('Axioms/KLE001+0.ax')	
include('Axioms/KLE001+4.ax')	
include('Axioms/KLE001+6.ax')	
$\forall x_0, x_1: (x_0 + x_1 = x_1 \Rightarrow \forall x_2: \text{forward_box}(x_0, \text{dom}(x_2)) + \text{forward_box}(x_1, \text{dom}(x_2)) = \text{forward_box}(x_0, \text{dom}(x_2)))$	fof(goals, conjecture)
KLE119+1.p Validity of abort rule	
The abort rule of Hoare logic is valid with respect to the Kleene algebra semantics.	
include('Axioms/KLE001+0.ax')	
include('Axioms/KLE001+4.ax')	
include('Axioms/KLE001+6.ax')	
$\forall x_0, x_1: \text{backward_diamond}(0, \text{dom}(x_0)) + \text{dom}(x_1) = \text{dom}(x_1)$	fof(goals, conjecture)
KLE120+1.p Validity of skip rule	
The skip rule of Hoare logic is valid with respect to the Kleene algebra semantics.	
include('Axioms/KLE001+0.ax')	

```

include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0: \text{backward\_diamond}(1, \text{dom}(x_0)) + \text{dom}(x_0) = \text{dom}(x_0)$  fof(goals, conjecture)

```

KLE121+1.p Validity of composition rule

The composition rule of Hoare logic is valid with respect to the Kleene algebra semantics.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1, x_2, x_3, x_4: ((\text{backward\_diamond}(x_3, \text{dom}(x_0)) + \text{dom}(x_1) = \text{dom}(x_1) \text{ and } \text{backward\_diamond}(x_4, \text{dom}(x_1)) + \text{dom}(x_2) = \text{dom}(x_2)) \Rightarrow \text{backward\_diamond}(x_3 \cdot x_4, \text{dom}(x_0)) + \text{dom}(x_2) = \text{dom}(x_2)$  fof(goals, conjecture)

```

KLE122+1.p Validity of conditional rule

The conditional rule of Hoare logic is valid with respect to the Kleene algebra semantics.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1, x_2: \text{if\_then\_else}(x_0, x_1, x_2) = \text{dom}(x_0) \cdot x_1 + \text{ad}(x_0) \cdot x_2$  fof(if_then_else_definition, axiom)
 $\forall x_0, x_1, x_2, x_3, x_4: ((\text{backward\_diamond}(x_0, \text{dom}(x_2) \cdot \text{dom}(x_3)) + \text{dom}(x_4) = \text{dom}(x_4) \text{ and } \text{backward\_diamond}(x_1, \text{ad}(x_2) \cdot \text{dom}(x_3)) + \text{dom}(x_4) = \text{dom}(x_4)) \Rightarrow \text{backward\_diamond}(\text{if\_then\_else}(x_2, x_0, x_1), \text{dom}(x_3)) + \text{dom}(x_4) = \text{dom}(x_4)$  fof(goals, conjecture)

```

KLE123+1.p Validity of while rule

The while rule of Hoare logic is valid with respect to the Kleene algebra semantics.

```

include('Axioms/KLE002+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1: \text{while\_do}(x_1, x_0) = (\text{dom}(x_1) \cdot x_0)^* \cdot \text{ad}(x_1)$  fof(while_do_definition, axiom)
 $\forall x_0, x_1, x_2: (\text{backward\_diamond}(x_0, \text{dom}(x_1) \cdot \text{dom}(x_2)) + \text{dom}(x_2) = \text{dom}(x_2) \Rightarrow \text{backward\_diamond}(\text{while\_do}(x_1, x_0), \text{dom}(x_2) \cdot \text{dom}(x_2) = \text{dom}(x_2))$  fof(goals, conjecture)

```

KLE124+1.p Validity of weakening rule

The weakening rule of Hoare logic is valid with respect to the Kleene algebra semantics.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
 $\forall x_0, x_1, x_2, x_3, x_4: ((\text{dom}(x_2) + \text{dom}(x_1) = \text{dom}(x_1) \text{ and } \text{backward\_diamond}(x_0, \text{dom}(x_1)) + \text{dom}(x_3) = \text{dom}(x_3) \text{ and } \text{dom}(x_3) + \text{dom}(x_4) = \text{dom}(x_4)) \Rightarrow \text{backward\_diamond}(x_0, \text{dom}(x_2)) + \text{dom}(x_4) = \text{dom}(x_4)$  fof(goals, conjecture)

```

KLE125+1.p Quasicommutation theorem

If x quasicommutes over y , then $x+y$ terminates if x and y individually do.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
include('Axioms/KLE001+7.ax')
 $\forall x_0, x_1: (x_0 \cdot x_1 + x_1 \cdot (x_1 + x_0)^* = x_1 \cdot (x_1 + x_0)^* \Rightarrow (\text{divergence}(x_1 + x_0) = 0 \iff \text{divergence}(x_1) + \text{divergence}(x_0) = 0))$  fof(goals, conjecture)

```

KLE126+1.p Lazy commutation theorem

If x lazily commutes over y , then $x+y$ terminates if x and y individually do.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
include('Axioms/KLE001+7.ax')
 $\forall x_0, x_1: (x_0 \cdot x_1 + x_1 \cdot (x_1 + x_0)^* = x_1 \cdot (x_1 + x_0)^* + x_0 \Rightarrow (\text{divergence}(x_1 + x_0) = 0 \iff \text{divergence}(x_1) + \text{divergence}(x_0) = 0))$  fof(goals, conjecture)

```

KLE127+1.p Loop refinement theorem

If x quasicommutes over y , then all infinite behaviours of $x+y$ can be separated into infinite behaviours of x and a infinite behaviour of y after finitely many x steps.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
include('Axioms/KLE001+7.ax')

```

$\forall x_0, x_1: (x_0 \cdot x_1 + x_1 \cdot (x_1 + x_0)^*) = x_1 \cdot (x_1 + x_0)^* \Rightarrow \text{divergence}(x_1 + x_0) = \text{divergence}(x_1) + \text{forward_diamond}(x_1^*, \text{divergence}(x_0))$

KLE128+1.p Comparison of two different notions of termination

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

include('Axioms/KLE001+7.ax')

$\forall x_0: (\text{divergence}(x_0) = 0 \Rightarrow \forall x_1: (\text{dom}(x_1) + \text{forward_diamond}(x_0, \text{dom}(x_1))) = \text{forward_diamond}(x_0, \text{dom}(x_1)) \Rightarrow \text{dom}(x_1) = 0)) \quad \text{fof(goals, conjecture)}$

KLE129+1.p Comparison of two different notions of termination

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

include('Axioms/KLE001+7.ax')

$\forall x_0: (\text{divergence}(x_0) = 0 \Leftarrow \forall x_1: (\text{dom}(x_1) + \text{forward_diamond}(x_0, \text{dom}(x_1))) = \text{forward_diamond}(x_0, \text{dom}(x_1)) \Rightarrow \text{dom}(x_1) = 0)) \quad \text{fof(goals, conjecture)}$

KLE130+1.p Two notions of termination

The lemma compares absence of divergence with a third notion of termination.

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

include('Axioms/KLE001+7.ax')

$\forall x_0: (\text{divergence}(x_0) = 0 \Rightarrow \forall x_1: \text{dom}(x_1) + \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{dom}(x_1), \text{forward_diamond}(x_0, \text{dom}(x_1)))) = \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{dom}(x_1), \text{forward_diamond}(x_0, \text{dom}(x_1)))))) \quad \text{fof(goals, conjecture)}$

KLE131+1.p Two notions of termination

The lemma compares absence of divergence with a third notion of termination.

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

include('Axioms/KLE001+7.ax')

$\forall x_0: (\text{divergence}(x_0) = 0 \Leftarrow \forall x_1: \text{dom}(x_1) + \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{dom}(x_1), \text{forward_diamond}(x_0, \text{dom}(x_1)))) = \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{dom}(x_1), \text{forward_diamond}(x_0, \text{dom}(x_1)))))) \quad \text{fof(goals, conjecture)}$

KLE132+1.p Every element that satisfies Loeb's formula is wellfounded

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

include('Axioms/KLE001+7.ax')

$\forall x_0: (\forall x_1: \text{forward_diamond}(x_0, \text{dom}(x_1)) + \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{dom}(x_1), \text{forward_diamond}(x_0, \text{dom}(x_1)))) = \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{dom}(x_1), \text{forward_diamond}(x_0, \text{dom}(x_1)))) \Rightarrow \forall x_2: (\text{dom}(x_2) + \text{forward_diamond}(x_0, \text{dom}(x_2))) = \text{forward_diamond}(x_0, \text{dom}(x_2)) \Rightarrow \text{dom}(x_2) = 0)) \quad \text{fof(goals, conjecture)}$

KLE133+1.p Loeb's formula and wellfoundedness

Every diamond transitive wellfounded element satisfies Loeb's formula.

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

include('Axioms/KLE001+7.ax')

$\forall x_0: ((\forall x_1: \text{dom}(x_1) + \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{dom}(x_1), \text{forward_diamond}(x_0, \text{dom}(x_1)))) = \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{dom}(x_1), \text{forward_diamond}(x_0, \text{dom}(x_1)))) \Rightarrow \forall x_2: \text{dom}(x_2) + \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{dom}(x_2), \text{forward_diamond}(x_0, \text{dom}(x_2)))) = \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{dom}(x_2), \text{forward_diamond}(x_0, \text{dom}(x_2)))) \Rightarrow \forall x_3: \text{dom}(x_3) + \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{dom}(x_3), \text{forward_diamond}(x_0, \text{dom}(x_3)))) = \text{forward_diamond}(x_0^*, \text{domain_difference}(\text{dom}(x_3), \text{forward_diamond}(x_0, \text{dom}(x_3)))))) \quad \text{fof(goals, conjecture)}$

KLE134+1.p A no name lemma

include('Axioms/KLE001+0.ax')

include('Axioms/KLE001+4.ax')

include('Axioms/KLE001+6.ax')

$\forall x_0, x_1, x_2: \text{domain_difference}(\text{forward_diamond}(x_0, \text{dom}(x_1)), \text{forward_diamond}(x_0, \text{dom}(x_2))) + \text{forward_diamond}(x_0, \text{domain_difference}(\text{forward_diamond}(x_0, \text{dom}(x_1)), \text{forward_diamond}(x_0, \text{dom}(x_2)))) = \text{forward_diamond}(x_0, \text{domain_difference}(\text{forward_diamond}(x_0, \text{dom}(x_1)), \text{forward_diamond}(x_0, \text{dom}(x_2)))) \quad \text{fof(goals, conjecture)}$

KLE135+1.p Two notions of termination

If an element does not diverge, then it always becomes disabled after finitely many steps.

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
include('Axioms/KLE001+7.ax')
 $\forall x_0: (\text{divergence}(x_0) = 0 \Rightarrow \text{forward\_diamond}(x_0^*, \text{ad}(x_0)) = 1) \quad \text{fof(goals, conjecture)}$ 

```

KLE136+1.p Newman's lemma holds in divergence Kleene algebras

```

include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
include('Axioms/KLE001+7.ax')
 $\forall x_0, x_1: ((\text{divergence}(x_0 + x_1) = 0 \text{ and } x_1 \cdot x_0 + x_0^* \cdot x_1^* = x_0^* \cdot x_1^*) \Rightarrow (x_0 + x_1)^* + x_0^* \cdot x_1^* = x_0^* \cdot x_1^*) \quad \text{fof(goals, conjecture)}$ 

```

KLE137+1.p There is a greatest element, namely 1 \wedge infty

```

include('Axioms/KLE004+0.ax')
 $\forall x_0: x_0 \leq \text{strong\_iteration}(1) \quad \text{fof(goals, conjecture)}$ 

```

KLE138+1.p Strong iteration of a abort is miracle

```

include('Axioms/KLE004+0.ax')
 $\text{strong\_iteration}(0) = 1 \quad \text{fof(goals, conjecture)}$ 

```

KLE139+1.p Dual unfold

```

include('Axioms/KLE004+0.ax')
 $\forall x_0: \text{strong\_iteration}(x_0) = \text{strong\_iteration}(x_0) \cdot x_0 + 1 \quad \text{fof(goals, conjecture)}$ 

```

KLE139+2.p Dual unfold

```

include('Axioms/KLE004+0.ax')
 $\forall x_0: (\text{strong\_iteration}(x_0) \leq \text{strong\_iteration}(x_0) \cdot x_0 + 1 \text{ and } \text{strong\_iteration}(x_0) \cdot x_0 + 1 \leq \text{strong\_iteration}(x_0)) \quad \text{fof(goals, conjecture)}$ 

```

KLE140+1.p Isotonicity of strong iteration

```

include('Axioms/KLE004+0.ax')
 $\forall x_0, x_1: (x_0 \leq x_1 \Rightarrow \text{strong\_iteration}(x_0) \leq \text{strong\_iteration}(x_1)) \quad \text{fof(goals, conjecture)}$ 

```

KLE141+1.p The greatest is left annihilator

The greatest is left annihilator; a miracle followed by anything is again a miracle.

```

include('Axioms/KLE004+0.ax')
 $\forall x_0: \text{strong\_iteration}(1) \cdot x_0 = \text{strong\_iteration}(1) \quad \text{fof(goals, conjecture)}$ 

```

KLE141+2.p The greatest is left annihilator

The greatest is left annihilator; a miracle followed by anything is again a miracle.

```

include('Axioms/KLE004+0.ax')
 $\forall x_0: (\text{strong\_iteration}(1) \cdot x_0 \leq \text{strong\_iteration}(1) \text{ and } \text{strong\_iteration}(1) \leq \text{strong\_iteration}(1) \cdot x_0) \quad \text{fof(goals, conjecture)}$ 

```

KLE142+1.p If strong iteration is applied twice, a miracle occurs.

```

include('Axioms/KLE004+0.ax')
 $\forall x_0: \text{strong\_iteration}(\text{strong\_iteration}(x_0)) = \text{strong\_iteration}(1) \quad \text{fof(goals, conjecture)}$ 

```

KLE142+2.p If strong iteration is applied twice, a miracle occurs.

```

include('Axioms/KLE004+0.ax')
 $\forall x_0: (\text{strong\_iteration}(\text{strong\_iteration}(x_0)) \leq \text{strong\_iteration}(1) \text{ and } \text{strong\_iteration}(1) \leq \text{strong\_iteration}(\text{strong\_iteration}(x_0))) \quad \text{fof(goals, conjecture)}$ 

```

KLE143+1.p Strong iteration is idempotent w.r.t. multiplication

```

include('Axioms/KLE004+0.ax')
 $\forall x_0: \text{strong\_iteration}(x_0) \cdot \text{strong\_iteration}(x_0) = \text{strong\_iteration}(x_0) \quad \text{fof(goals, conjecture)}$ 

```

KLE143+2.p Strong iteration is idempotent w.r.t. multiplication

```

include('Axioms/KLE004+0.ax')
 $\forall x_0: (\text{strong\_iteration}(x_0) \cdot \text{strong\_iteration}(x_0) \leq \text{strong\_iteration}(x_0) \text{ and } \text{strong\_iteration}(x_0) \leq \text{strong\_iteration}(x_0) \cdot \text{strong\_iteration}(x_0)) \quad \text{fof(goals, conjecture)}$ 

```

KLE144+1.p Strong iteration applied after finite iteration is magic.

```

include('Axioms/KLE004+0.ax')
 $\forall x_0: \text{strong\_iteration}(x_0^*) = \text{strong\_iteration}(1) \quad \text{fof(goals, conjecture)}$ 

```

KLE144+2.p Strong iteration applied after finite iteration is magic.

```

include('Axioms/KLE004+0.ax')
 $\forall x_0: (\text{strong\_iteration}(x_0^*) \leq \text{strong\_iteration}(1) \text{ and } \text{strong\_iteration}(1) \leq \text{strong\_iteration}(x_0^*)) \quad \text{fof(goals, conjecture)}$ 

```

KLE145+1.p Finite iteration after strong iteration is strong iteration

```
include('Axioms/KLE004+0.ax')
 $\forall x_0: \text{strong\_iteration}(x_0)^* = \text{strong\_iteration}(x_0)$  fof(goals, conjecture)
```

KLE145+2.p Finite iteration after strong iteration is strong iteration

```
include('Axioms/KLE004+0.ax')
 $\forall x_0: (\text{strong\_iteration}(x_0)^* \leq \text{strong\_iteration}(x_0) \text{ and } \text{strong\_iteration}(x_0) \leq \text{strong\_iteration}(x_0)^*)$  fof(goals, conjecture)
```

KLE146+1.p Skip is part of strong iteration

```
include('Axioms/KLE004+0.ax')
 $\forall x_0: 1 \leq \text{strong\_iteration}(x_0)$  fof(goals, conjecture)
```

KLE147+1.p Sliding of strong iteration

```
include('Axioms/KLE004+0.ax')
 $\forall x_0, x_1: \text{strong\_iteration}(x_0^* \cdot x_1) = x_1^* \cdot \text{strong\_iteration}(x_0^* \cdot x_1)$  fof(goals, conjecture)
```

KLE147+2.p Sliding of strong iteration

```
include('Axioms/KLE004+0.ax')
 $\forall x_0, x_1: (\text{strong\_iteration}(x_0^* \cdot x_1) \leq x_1^* \cdot \text{strong\_iteration}(x_0^* \cdot x_1) \text{ and } x_1^* \cdot \text{strong\_iteration}(x_0^* \cdot x_1) \leq \text{strong\_iteration}(x_0^* \cdot x_1))$  fof(goals, conjecture)
```

KLE148+1.p Blocking law

If y is blocked by x then x before strong iteration of y reduces to x.

```
include('Axioms/KLE004+0.ax')
 $\forall x_0, x_1: (x_0 \cdot x_1 = 0 \Rightarrow x_0 \cdot \text{strong\_iteration}(x_1) = x_0)$  fof(goals, conjecture)
```

KLE148+2.p Blocking law

If y is blocked by x then x before strong iteration of y reduces to x.

```
include('Axioms/KLE004+0.ax')
 $\forall x_0, x_1: ((x_0 \cdot x_1 = 0 \Rightarrow x_0 \cdot \text{strong\_iteration}(x_1) \leq x_0) \text{ and } x_0 \leq x_0 \cdot \text{strong\_iteration}(x_1))$  fof(goals, conjecture)
```

KLE149+1.p Strong version of unfold

```
include('Axioms/KLE004+0.ax')
 $\forall x_0, x_1: \text{strong\_iteration}(x_0 + x_1) = (x_1^* \cdot x_0) \cdot \text{strong\_iteration}(x_0 + x_1) + \text{strong\_iteration}(x_1)$  fof(goals, conjecture)
```

KLE149+2.p Strong version of unfold

```
include('Axioms/KLE004+0.ax')
 $\forall x_0, x_1: (\text{strong\_iteration}(x_0 + x_1) \leq (x_1^* \cdot x_0) \cdot \text{strong\_iteration}(x_0 + x_1) + \text{strong\_iteration}(x_1) \text{ and } (x_1^* \cdot x_0) \cdot \text{strong\_iteration}(x_0 + x_1) + \text{strong\_iteration}(x_1) \leq \text{strong\_iteration}(x_0 + x_1))$  fof(goals, conjecture)
```

KLE150+1.p Iterating non-terminating elements reduces to the element itself

```
include('Axioms/KLE004+0.ax')
 $\forall x_0: \text{strong\_iteration}(x_0 \cdot 0) = 1 + x_0 \cdot 0$  fof(goals, conjecture)
```

KLE150+2.p Iterating non-terminating elements reduces to the element itself

```
include('Axioms/KLE004+0.ax')
 $\forall x_0: (\text{strong\_iteration}(x_0 \cdot 0) \leq 1 + x_0 \cdot 0 \text{ and } 1 + x_0 \cdot 0 \leq \text{strong\_iteration}(x_0 \cdot 0))$  fof(goals, conjecture)
```

KLE151+1.p Sliding of strong iteration

Two ways of grouping an alternation of x's and y's that starts and ends with x.

```
include('Axioms/KLE004+0.ax')
 $\forall x_0, x_1: x_0 \cdot \text{strong\_iteration}(x_1 \cdot x_0) \leq \text{strong\_iteration}(x_0 \cdot x_1) \cdot x_0$  fof(goals, conjecture)
```

KLE152+1.p Sliding of strong iteration

Two ways of grouping an alternation of x's and y's that starts and ends with x.

```
include('Axioms/KLE004+0.ax')
 $\forall x_0, x_1: \text{strong\_iteration}(x_0 \cdot x_1) \cdot x_0 \leq x_0 \cdot \text{strong\_iteration}(x_1 \cdot x_0)$  fof(goals, conjecture)
```

KLE153+1.p Sliding of strong iteration

Two ways of grouping an alternation of x's and y's that starts and ends with x.

```
include('Axioms/KLE004+0.ax')
 $\forall x_0, x_1: x_0 \cdot \text{strong\_iteration}(x_1 \cdot x_0) = \text{strong\_iteration}(x_0 \cdot x_1) \cdot x_0$  fof(goals, conjecture)
```

KLE154+1.p Denesting of strong iteration

```
include('Axioms/KLE004+0.ax')
 $\forall x_0, x_1: \text{strong\_iteration}(x_0 + x_1) = \text{strong\_iteration}(x_0) \cdot \text{strong\_iteration}(x_1 \cdot \text{strong\_iteration}(x_0))$  fof(goals, conjecture)
```

KLE154+2.p Denesting of strong iteration

```
include('Axioms/KLE004+0.ax')
```

$\forall x_0, x_1: (\text{strong_iteration}(x_0 + x_1) \leq \text{strong_iteration}(x_0) \cdot \text{strong_iteration}(x_1) \cdot \text{strong_iteration}(x_0)) \text{ and } \text{strong_iteration}(x_0) \cdot \text{strong_iteration}(x_1 \cdot \text{strong_iteration}(x_0)) \leq \text{strong_iteration}(x_0 + x_1)) \quad \text{fof(goals, conjecture)}$

KLE155+1.p Denesting of strong iteration

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1: \text{strong_iteration}(x_0^* \cdot x_1) \cdot \text{strong_iteration}(x_0) = \text{strong_iteration}(x_0 + x_1) \quad \text{fof(goals, conjecture)}$

KLE155+2.p Denesting of strong iteration

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1: (\text{strong_iteration}(x_0 + x_1) \leq \text{strong_iteration}(x_0^* \cdot x_1) \cdot \text{strong_iteration}(x_0) \text{ and } \text{strong_iteration}(x_0^* \cdot x_1) \cdot \text{strong_iteration}(x_0) \leq \text{strong_iteration}(x_0 + x_1)) \quad \text{fof(goals, conjecture)}$

KLE156+1.p Semicommuation law of finite iteration

If x semicomutes over y every finite sequence of x's and y's can be rearranged to a finite sequence of x's followed by finite sequence of y's.

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1: (x_0 \cdot x_1 \leq x_1 \cdot x_0 \Rightarrow (x_1 + x_0)^* = x_1^* \cdot x_0^*) \quad \text{fof(goals, conjecture)}$

KLE156+2.p Semicommuation law of finite iteration

If x semicomutes over y every finite sequence of x's and y's can be rearranged to a finite sequence of x's followed by finite sequence of y's.

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1: ((x_0 \cdot x_1 \leq x_1 \cdot x_0 \Rightarrow (x_1 + x_0)^* \leq x_1^* \cdot x_0^*) \text{ and } x_1^* \cdot x_0^* \leq (x_1 + x_0)^*) \quad \text{fof(goals, conjecture)}$

KLE157+1.p Semicommuation law of finite iteration

If x semicomutes over y every sequence of x's and y's can be rearranged to a sequence of x's followed by sequence of y's.

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1: (x_0 \cdot x_1 \leq x_1 \cdot x_0 \Rightarrow \text{strong_iteration}(x_1 + x_0) = \text{strong_iteration}(x_1) \cdot \text{strong_iteration}(x_0)) \quad \text{fof(goals, conjecture)}$

KLE157+2.p Semicommuation law of finite iteration

If x semicomutes over y every sequence of x's and y's can be rearranged to a sequence of x's followed by sequence of y's.

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1: ((x_0 \cdot x_1 \leq x_1 \cdot x_0 \Rightarrow \text{strong_iteration}(x_1 + x_0) \leq \text{strong_iteration}(x_1) \cdot \text{strong_iteration}(x_0)) \text{ and } \text{strong_iteration}(x_0) \leq \text{strong_iteration}(x_1 + x_0)) \quad \text{fof(goals, conjecture)}$

KLE158+1.p Simulation law for data refinement

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1, x_2: (x_0 \cdot x_1 \leq x_2 \cdot x_0 \Rightarrow x_0 \cdot \text{strong_iteration}(x_1) \leq \text{strong_iteration}(x_2) \cdot x_0) \quad \text{fof(goals, conjecture)}$

KLE159+1.p Simulation law for data refinement

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1, x_2: (x_0 \cdot x_1 \leq x_2 \cdot x_0 \Rightarrow x_0 \cdot x_1^* \leq x_2^* \cdot x_0) \quad \text{fof(goals, conjecture)}$

KLE160+1.p Simulation law for data refinement

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1, x_2: (x_0 \cdot x_1 \leq x_1 \cdot x_2 \Rightarrow x_0^* \cdot x_1 \leq x_1 \cdot x_2^*) \quad \text{fof(goals, conjecture)}$

KLE161+1.p Data refinement

The first hypothesis says that b cannot loop infinitely. The second hypothesis says that a is data refined by aa w.r.t. upward simulation z. By the third hypothesis, 1 is data refined by b. The fourth and fifth condition expresses the standard data refinement of initialisations and finalisations.

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1, x_2, x_3, x_4, x_5, x_6, x_7: ((\text{strong_iteration}(x_7) = x_7^* \text{ and } x_0 \cdot x_6 \leq x_5 \cdot x_0 \text{ and } x_0 \cdot x_7 \leq x_0 \text{ and } x_2 \leq x_1 \cdot x_0 \text{ and } x_0 \cdot x_4 \leq x_3) \Rightarrow x_2 \cdot (\text{strong_iteration}(x_6 + x_7) \cdot x_4) \leq x_1 \cdot (\text{strong_iteration}(x_5) \cdot x_3)) \quad \text{fof(goals, conjecture)}$

KLE162+1.p Part 1 of Back's atomicity refinement theorem

Back's atomicity refinement theorem is proved up to the reconstruction of concurrency.

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1, x_2, x_3, x_4, x_5: ((x_0 \leq x_3 \cdot x_0 \text{ and } x_4 \cdot x_1 \leq x_1 \cdot x_4 \text{ and } (x_0 + (x_4 + x_1)) \cdot x_2 \leq x_2 \cdot (x_0 + (x_4 + x_1)) \text{ and } x_4 \cdot x_3 \leq x_3 \cdot x_4 \text{ and } x_4^* = \text{strong_iteration}(x_4) \text{ and } x_3 \cdot x_2 \leq x_2 \cdot x_3 \text{ and } x_3 \cdot x_1 = 0 \text{ and } x_5 \leq x_5 \cdot x_3) \Rightarrow x_5 \cdot (\text{strong_iteration}((x_0 + (x_4 + x_1)) + x_2) \cdot x_3) \leq (x_5 \cdot (\text{strong_iteration}(x_2) \cdot x_3)) \cdot ((\text{strong_iteration}(x_4) \cdot x_3) \cdot \text{strong_iteration}((x_0 \cdot \text{strong_iteration}(x_1)) \cdot (x_3 \cdot \text{strong_iteration}(x_4)))) \quad \text{fof(goals, conjecture)}$

KLE163+1.p Part 2 of Back's atomicity refinement theorem

The concurrency bit of Back's atomicity refinement theorem.

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1, x_2, x_3, x_4, x_5: ((x_0 \leq x_3 \cdot x_0 \text{ and } x_4 \cdot x_1 \leq x_1 \cdot x_4 \text{ and } (x_0 + (x_4 + x_1)) \cdot x_2 \leq x_2 \cdot (x_0 + (x_4 + x_1)) \text{ and } x_4 \cdot x_3 \leq x_3 \cdot x_4 \text{ and } x_4^* = \text{strong_iteration}(x_4) \text{ and } x_3 \cdot x_2 \leq x_2 \cdot x_3 \text{ and } x_3 \cdot x_1 = 0 \text{ and } x_5 \leq x_5 \cdot x_3 \text{ and } x_3 \leq 1) \Rightarrow (x_5 \cdot (\text{strong_iteration}(x_2) \cdot x_3)) \cdot ((\text{strong_iteration}(x_4) \cdot x_3) \cdot \text{strong_iteration}((x_0 \cdot \text{strong_iteration}(x_1)) \cdot (x_3 \cdot \text{strong_iteration}(x_4)))) \leq x_5 \cdot \text{strong_iteration}(((x_0 \cdot \text{strong_iteration}(x_1)) \cdot x_3 + x_4) + x_2))$ fof(goals, conjecture)

KLE164+1.p Back's atomicity refinement theorem

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1, x_2, x_3, x_4, x_5: ((x_0 \leq x_3 \cdot x_0 \text{ and } x_4 \cdot x_1 \leq x_1 \cdot x_4 \text{ and } (x_0 + (x_4 + x_1)) \cdot x_2 \leq x_2 \cdot (x_0 + (x_4 + x_1)) \text{ and } x_4 \cdot x_3 \leq x_3 \cdot x_4 \text{ and } x_4^* = \text{strong_iteration}(x_4) \text{ and } x_3 \cdot x_2 \leq x_2 \cdot x_3 \text{ and } x_3 \cdot x_1 = 0 \text{ and } x_5 \leq x_5 \cdot x_3 \text{ and } x_3 \leq 1) \Rightarrow x_5 \cdot (\text{strong_iteration}((x_0 + (x_4 + x_1)) + x_2) \cdot x_3) \leq x_5 \cdot \text{strong_iteration}(((x_0 \cdot \text{strong_iteration}(x_1)) \cdot x_3 + x_4) + x_2))$ fof(goals, conjecture)

KLE165+1.p Denest for weakly quasi commutation

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1: (x_0 \cdot x_1 \leq x_1 \cdot \text{strong_iteration}(x_1 + x_0) \Rightarrow \text{strong_iteration}(x_1 + x_0) = \text{strong_iteration}(x_1) \cdot \text{strong_iteration}(x_0))$ fof(goals, conjecture)

KLE165+2.p Denest for weakly quasi commutation

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1: ((x_0 \cdot x_1 \leq x_1 \cdot \text{strong_iteration}(x_1 + x_0) \Rightarrow \text{strong_iteration}(x_1 + x_0) \leq \text{strong_iteration}(x_1) \cdot \text{strong_iteration}(x_0)) \text{ and } \text{strong_iteration}(x_0) \leq \text{strong_iteration}(x_1 + x_0))$ fof(goals, conjecture)

KLE166+1.p Strong iteration

If x weakly quasicommutates over y, x and y is strongly terminating iff x + y strongly terminates.

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1: ((x_0 \cdot x_1 \leq x_1 \cdot \text{strong_iteration}(x_1 + x_0) \Rightarrow (\text{strong_iteration}(x_1 + x_0) \cdot 0 = 0 \iff \text{strong_iteration}(x_1) \cdot 0 + \text{strong_iteration}(x_0) \cdot 0 = 0))$ fof(goals, conjecture)

KLE167+1.p Blocking law

If y is blocked by x then x before finite iteration of y reduces to x.

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1: ((x_0 \cdot x_1 = 0 \Rightarrow x_0 \cdot x_1^* \leq x_0) \text{ and } x_0 \leq x_0 \cdot x_1^*)$ fof(goals, conjecture)

KLE168+1.p If x quasicommutates over y then x \wedge infty quasicommutates over y \wedge infty

include('Axioms/KLE004+0.ax')

$\forall x_0, x_1: (x_0 \cdot x_1 \leq x_1 \cdot x_0 \Rightarrow \text{strong_iteration}(x_0) \cdot \text{strong_iteration}(x_1) \leq \text{strong_iteration}(x_1) \cdot \text{strong_iteration}(x_0))$ fof(goals, conjecture)

KLE169+1.p Exponential automata

Automata grows exponentially with N in sigma*.a.sigma \wedge N.a

include('Axioms/KLE002+0.ax')

sigma = a + b fof(an, axiom)

$a \cdot (b \cdot a) \leq \text{sigma}^* \cdot (a \cdot (\text{sigma} \cdot a))$ fof(a, conjecture)

KLE170+1.002.p a \wedge N \leq a*, N=2

include('Axioms/KLE002+0.ax')

$a \cdot a \leq a^*$ fof(a, conjecture)

KLE170+1.004.p a \wedge N \leq a*, N=4

include('Axioms/KLE002+0.ax')

$a \cdot (a \cdot (a \cdot a)) \leq a^*$ fof(a, conjecture)

KLE170+1.006.p a \wedge N \leq a*, N=6

include('Axioms/KLE002+0.ax')

$a \cdot (a \cdot (a \cdot (a \cdot a))) \leq a^*$ fof(a, conjecture)

KLE171+1.p Ben's problem 1

include('Axioms/KLE002+0.ax')

sigma = a + b fof(an, axiom)

$a \cdot (b \cdot (b \cdot a)) \leq \text{sigma}^* \cdot (a \cdot (\text{sigma} \cdot a))$ fof(a, conjecture)

KLE172+1.p Ben's problem 2

include('Axioms/KLE002+0.ax')

sigma = a + b fof(an, axiom)

$a \cdot (b \cdot (b \cdot (a \cdot b))) \leq \text{sigma}^* \cdot (a \cdot (\text{sigma} \cdot a))$ fof(a, conjecture)

KLE173+1.p Idempotent semirings with tests

include('Axioms/KLE001+0.ax')

```
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
include('Axioms/KLE001+3.ax')
```

KLE174+1.p Idempotent semirings with domain/codomain, modal

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
include('Axioms/KLE001+7.ax')
```

KLE175+1.p Idempotent semirings with tests

```
include('Axioms/KLE001+0.ax')
include('Axioms/KLE001+5.ax')
```

KLE176+1.p Kleene algebra

```
include('Axioms/KLE002+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
include('Axioms/KLE001+3.ax')
```

KLE177+1.p Kleene algebra

```
include('Axioms/KLE002+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
include('Axioms/KLE001+7.ax')
```

KLE178+1.p Kleene Algebra

```
include('Axioms/KLE002+0.ax')
include('Axioms/KLE001+5.ax')
```

KLE179+1.p Omega algebra

```
include('Axioms/KLE003+0.ax')
include('Axioms/KLE001+1.ax')
include('Axioms/KLE001+2.ax')
include('Axioms/KLE001+3.ax')
```

KLE180+1.p Omega algebra

```
include('Axioms/KLE003+0.ax')
include('Axioms/KLE001+4.ax')
include('Axioms/KLE001+6.ax')
include('Axioms/KLE001+7.ax')
```

KLE181+1.p Omega algebra

```
include('Axioms/KLE003+0.ax')
include('Axioms/KLE001+5.ax')
```

KLE182+1.p Demonic Refinement Algebra

```
include('Axioms/KLE004+0.ax')
```